

From flows of Λ Fleming-Viot processes to lookdown processes via flows of partitions

Cyril Labbé

Laboratoire de Probabilités et Modèles Aléatoires

Université Pierre et Marie Curie

cyril.labbe@upmc.fr

—

February 13, 2012

Abstract

We unify two well-known representations of Λ Fleming-Viot processes: the stochastic flow of bridges and the lookdown process. We show that the stochastic flow of bridges can be uniquely decomposed into two new objects: the Eves process, which are the genetic types that play a significant rôle in the population as time passes, and a stochastic flow of partitions, which encodes the genealogical structure. We prove that the latter is completely equivalent with the so-called lookdown graph and so, we provide a pathwise connection between flows of bridges and lookdown processes: using the lookdown construction with that flow of partitions and the Eves, we recover pathwise the collection of Λ Fleming-Viot processes encoded by the flow of bridges.

1 Introduction

A generalized Fleming-Viot process $\rho := (\rho_t, t \geq 0)$ is a Markov process that describes the evolution of an infinite population. It takes its values in the set of probability measures on $[0, 1]$, where each point in $[0, 1]$ should be understood as a *genetic type*. For any $a \leq b \in [0, 1]$, $\rho_t([a, b])$ is the proportion of individuals at time $t \geq 0$ who descend from ancestors with types in $[a, b]$ and thus, ρ_t describes the composition of the population at time t . Bertoin and Le Gall in [6] show that the distribution of a generalized Fleming-Viot process is completely characterized by a finite measure Λ on $[0, 1]$, thus ρ is also called a Λ Fleming-Viot process. This process starts from ρ_0 taken as the uniform distribution on $[0, 1]$, and evolves through reproduction events that can be informally described as follows. At rate $x^{-2}\Lambda(dx)$ a parent is uniformly chosen among the population and gives its type to a fraction x of individuals, see Subsection

MSC 2010 subject classifications. Primary 60J25; secondary 60G09, 92D25.

Key words and phrases. Stochastic flow, Fleming-Viot process, Lookdown process, Partition-valued process, Coalescence, Continuous state branching process, Bridge

2.3 for a precise definition. Bertoin and Le Gall prove in [6] that there exists a random point $\mathbf{e}$, uniformly distributed over $[0, 1]$, such that

$$\rho_t(\{\mathbf{e}\}) \xrightarrow[t \uparrow \infty]{} 1 \quad (1)$$

In words, it means that a type fixes (possibly in infinite time) in the population. This type is called the *primitive Eve* of the population, it gives a first approximation of the composition of the population when t becomes large. It is natural to ask if finer results about the composition of the population can be obtained, such as the existence of a second Eve (in a sense to be made precise) for instance.

A point $x \in [0, 1]$ will be called an *ancestral type* if there exists $t > 0$ such that $\rho_t(\{x\}) > 0$. A simple argument (see Subsection 4.1) entails that the collection of ancestral types is countable. We will say that an ancestral type x *becomes extinct* if $d := \sup\{t > 0 : \rho_t(\{x\}) > 0\}$ is finite, d is then called its extinction time.

An important subclass of Λ Fleming-Viot processes are those who enjoy the following property. Almost surely, for every $t > 0$, ρ_t is a weighted sum of a finite number of Dirac masses on $[0, 1]$. Such Λ Fleming-Viot processes are said to come down from infinity (CDI), and then one can easily prove that the primitive Eve fixes in finite time. Roughly speaking, the (CDI) property means that on any interval of time $[0, \epsilon]$ infinitely many ancestral types become extinct, and the population at time ϵ is only composed of a finite number of types. This is the case when $\Lambda(dx) = \delta_0(dx)$ (standard Fleming-Viot process / Kingman coalescent) and when Λ is the density of a Beta($2 - \alpha, \alpha$) r.v. with $\alpha \in (1, 2)$ (Beta($2 - \alpha, \alpha$) Fleming-Viot process / Beta($2 - \alpha, \alpha$) coalescent), whereas when $\Lambda(dx) = dx$ (Neveu branching process / Bolthausen-Sznitman coalescent) it does not hold. When the Λ Fleming-Viot comes down from infinity, we define the following event

$$E := \{\text{There exists } t > 0 \text{ s.t. two ancestral types become extinct simultaneously at time } t\} \quad (2)$$

To the best of our knowledge, the question of simultaneous extinction of ancestral types has never been addressed.

Theorem 1 *The event E is trivial, that is, $\mathbb{P}(E) \in \{0, 1\}$. In addition, for a Beta($2 - \alpha, \alpha$) Fleming-Viot process with $\alpha \in (1, 2]$, $\mathbb{P}(E) = 0$.*

Note that for simplicity, the abusive notation Beta($0, 2$) (i.e. $\alpha = 2$) refers to the standard Fleming-Viot process. We conjecture that $\mathbb{P}(E) = 0$ for any Λ Fleming-Viot that comes down from infinity.

Suppose that the Λ Fleming-Viot process comes down from infinity and that $\mathbb{P}(E) = 0$. Its number of ancestral types is a non-increasing process that decreases by jumps of size 1. We define d^i as the jump time from i to $i - 1$ and $\mathbf{e}^i$ as the ancestral type that becomes extinct at this time, for every $i \geq 2$. In addition, let $\mathbf{e}^1$ be the ancestral type that fixes, that is, the primitive Eve. Thus, we obtain an ordering of the ancestral types $(\mathbf{e}^i)_{i \geq 1}$ by decreasing persistence: this gives precise information on the composition of the population as time passes and is consistent with the definition of the primitive Eve of Bertoin and Le Gall. This sequence is called the Eves.

When the Λ Fleming-Viot does not come down from infinity, none of the ancestral types become extinct so we need a different definition of the Eves. We restrict our construction to the Bolthausen-Sznitman case ($\Lambda(dx) = dx$) for a reason presented below and developed in Subsection 4.2. In the Bolthausen-Sznitman case, the primitive Eve fixes in *infinite* time : $\rho_t(\{e^1\}) \in (0, 1)$ for all $t > 0$. Proposition 4.7 shows that there exists a second Eve e^2 among $[0, 1] \setminus \{e^1\}$ that verifies

$$\lim_{t \rightarrow \infty} \frac{\rho_t(\{e^2\})}{1 - \rho_t(\{e^1\})} = 1$$

We then define recursively the complete sequence $(e^i)_{i \geq 1}$ following the same idea. One can wonder if this scheme could be applied to other measures Λ . To see that this question is not obvious, observe that for the Beta($2 - \alpha, \alpha$) Fleming-Viot processes with $\alpha \in (0, 1)$ there does not exist a second Eve.

So far, we have defined a sequence of Eves $(e^i)_{i \geq 1}$, which is an ordering of the ancestral types, in two different cases : **[(CDI) and $\mathbb{P}(E) = 0$]** and **[Bolthausen-Sznitman]**. In what follows, all the results hold indifferently in both cases. Proposition 4.11 proves that $(e^i)_{i \geq 1}$ is a sequence of i.i.d. uniform $[0, 1]$ r.v. and is independent of the sequence of processes $(\rho_t(\{e^i\}), t \geq 0)_{i \geq 1}$. This result means that the evolution of the asymptotic frequencies do not depend on the values taken by the Eves and is coherent with the fact that the Fleming-Viot model is neutral.

Let us turn our attention to a stationary version of this population model provided by the flow of bridges introduced by Bertoin and Le Gall in [6, 7, 8]. A flow of bridges $(F_{s,t}, -\infty < s \leq t < \infty)$ is a consistent collection of distribution functions verifying a property of independence and stationarity of its increments. Each distribution function $F_{s,t}$ defines a random probability measure $\rho_{s,t}$ on $[0, 1]$ such that the process $[s, \infty) \ni t \mapsto \rho_{s,t}$ is a Λ Fleming-Viot, for each $s \in \mathbb{R}$. Therefore $(\rho_{s,t}, t \in [s, \infty))_{s \in \mathbb{R}}$ is called a flow of Λ Fleming-Viot processes. Further details about this object will be provided in Subsection 2.2, but intuitively it should be seen as a stationary model of an infinite population where time is indexed by the real line. In this setting, $\rho_{s,t}([a, b])$ is the proportion of individuals alive at time t who descend from ancestors at time $s < t$ with types between a and b .

Using this flow representation, we can define for every $s \in \mathbb{R}$, the Eves $(e_s^i)_{i \geq 1}$ of the Λ Fleming-Viot process $(\rho_{s,t}, t \in [s, \infty))$. We call $(e_s^i, s \in \mathbb{R})_{i \geq 1}$ the *Eves process*. Of main interest is the question of the genealogical relationships between these Eves at distinct times. Fix $s < t$ and let $\hat{\Pi}_{s,t}$ be the random partition of the integers whose blocks, denoted $(\hat{\Pi}_{s,t}(i))_{i \geq 1}$ in the increasing order of their least element, are defined by

$$\hat{\Pi}_{s,t}(i) := \{j \in \mathbb{N} : e_t^j \text{ descends from } e_s^i\}$$

In words, we gather in a same block all the Eves at time t who descend from a same Eve at time s .

Theorem 2 *The collection of partitions $(\hat{\Pi}_{s,t}, -\infty < s \leq t < \infty)$ enjoys the following properties:*

- For every $r < s < t$, $\hat{\Pi}_{r,t} = \text{Coag}(\hat{\Pi}_{s,t}, \hat{\Pi}_{r,s})$ a.s. (cocycle property).
- $\hat{\Pi}_{s,t}$ is an exchangeable random partition whose law only depends on $t - s$. Furthermore, for any $s_1 < s_2 < \dots < s_n$ the partitions $\hat{\Pi}_{s_1, s_2}, \hat{\Pi}_{s_2, s_3}, \dots, \hat{\Pi}_{s_{n-1}, s_n}$ are independent.

- $\hat{\Pi}_{0,0} = 0_{[\infty]}$ and $\hat{\Pi}_{0,t} \rightarrow 0_{[\infty]}$ in probability as $t \downarrow 0$, for the distance $d_{\mathcal{P}}$.

Moreover the process $(\hat{\Pi}_{-t,0}, t \geq 0)$ is a Λ coalescent.

This collection is then called a stochastic flow of partitions associated with the measure Λ - or Λ flow of partitions for short. This object gives major information on the flow of Λ Fleming-Viot processes :

- Backward-in-time: the process $(\hat{\Pi}_{-t,0}, t \geq 0)$ gives the genealogy of the sequence of individuals $(\mathbf{e}_0^i)_{i \geq 1}$. In particular, $\hat{\Pi}_{-t,0}(j)$ is the subset, among the sampled individuals $(\mathbf{e}_0^i)_{i \geq 1}$, who found their common ancestor by time $-t$; this common ancestor having the type $\mathbf{e}_{-t}^j$.
- Forward-in-time: the process $(\hat{\Pi}_{0,t}, t \geq 0)$ gives the descendents of the Eves $(\mathbf{e}_0^i)_{i \geq 1}$ as time passes. In particular $\hat{\Pi}_{0,t}(i)$ is the collection of individuals alive at time $t > 0$ who descend from $\mathbf{e}_0^i$.

Notice that those two processes do not have the same distribution even if their one-dimensional marginals are equal in law. We make use of symbol "hat" in $\hat{\Pi}$ to emphasize the fact that the process goes forward-in-time, by opposition to the usual backward-in-time evolution of partition-valued processes that encode the genealogy.

We have identified pathwise from the flow of Λ Fleming-Viot two objects: the Eves process and a flow of partitions. The next theorem shows that they are sufficient to recover completely the flow of Λ Fleming-Viot. Introduce the notation $\mathcal{E}_s(\hat{\Pi}, (\mathbf{e}_s^i)_{i \geq 1})$ to designate the measure-valued process

$$[s, \infty) \ni t \mapsto \sum_{i=1}^{\infty} |\hat{\Pi}_{s,t}(i)| \delta_{\mathbf{e}_s^i}(dx)$$

where $|\hat{\Pi}_{s,t}(i)|$ stands for the asymptotic frequency of the i -th block of the exchangeable partition $\hat{\Pi}_{s,t}$.

Theorem 3 *The flow of Λ Fleming-Viot processes is uniquely decomposed into two random objects : the flow of partitions and the Eves process. More precisely, for each $s \in \mathbb{R}$ almost surely*

$$\mathcal{E}_s(\hat{\Pi}, (\mathbf{e}_s^i)_{i \geq 1}) = (\rho_{s,t}, t \in [s, \infty))$$

Furthermore if $\hat{\Pi}'$ is a Λ flow of partitions and for each $s \in \mathbb{R}$, $(\chi_s(i))_{i \geq 1}$ is a sequence of r.v. taking distinct values in $[0, 1]$ such that we have almost surely $\mathcal{E}_s(\hat{\Pi}', (\chi_s(i))_{i \geq 1}) = (\rho_{s,t}, t \in [s, \infty))$ then

- For each $s \in \mathbb{R}$, a.s. $(\chi_s(i))_{i \geq 1} = (\mathbf{e}_s^i)_{i \geq 1}$.
- Almost surely, $\hat{\Pi}' = \hat{\Pi}$.

This result also provides a striking connection with the lookdown representation of Λ Fleming-Viot processes due to Donnelly and Kurtz [11, 12]. Let us first recall briefly its definition. A lookdown process is a countable particle system whose process of limiting empirical measures is a Λ Fleming-Viot process. The particle system is entirely defined if we are given a sequence of initial types (one for each particle) and the dynamics of the particles. This dynamics is captured by the so-called lookdown graph (see for

instance [16]) that gives the genealogical relationships into the population, a heuristic definition of this object can be found at the end of this introduction. The connection with our decomposition theorem arises from the bijective correspondence between lockdown graphs and flows of partitions, which is proved in Section 3. Thanks to this connection, the flow of partitions $\hat{\Pi}$ appears as the unique lockdown graph that captures the genealogical relationships of the flow of bridges and we are able to define a *flow* of lockdown processes $(\xi_{s,t}(i), s \leq t)_{i \geq 1}$ by setting

- Initial types : $\xi_{s,s}(i) = \mathbf{e}_s^i$ for each $i \geq 1$.
- Dynamics : $\xi_{s,t}(j) = \xi_{s,s}(i) \Leftrightarrow j \in \hat{\Pi}_{s,t}(i)$

This sheds a new light on Theorem 3: $\mathcal{E}_s(\hat{\Pi}, (\mathbf{e}_s^i)_{i \geq 1})$ appears as the process of limiting empirical measures of the lockdown process $(\xi_{s,t}(i), t \in [s, \infty))_{i \geq 1}$ and is almost surely equal to $(\rho_{s,t}, t \in [s, \infty))$. So we obtain a coupling between the flow of bridges and the lockdown representation.

At first sight, it could seem simple to construct a flow of Λ Fleming-Viot from a lockdown graph using the lockdown construction: given a sequence of initial types at every time s we would obtain a collection of Λ Fleming-Viot processes by taking the limiting empirical measures of each lockdown process. However this collection does not form a flow of Λ Fleming-Viot unless the initial types were coupled so that they form the Eves process. This result is a direct consequence of Theorem 3.

A heuristic definition of a lockdown graph

A lockdown graph is a random point collection $\mathcal{P} = \{(t, u)\}$, where each point (t, u) is called a *reproduction event*: t is the time at which the reproduction occurs while u is a sequence of 0 and 1 that tells which levels are involved in the reproduction event (a level involved corresponds to the value 1 in the sequence). The lowest level involved is called the parent, it gives at time t its type to all the other levels involved. In Figure 1, this is represented by a collection of arrows: each of them starts from the level of the parent and hits a level involved. When an arrow hits a level, a new *line* is created at this level, while the previous line that occupied this level is pushed up to the next available level. From this picture, a collection of lines appears. The line located at level 1 is never pushed up while all the other lines only evolve by push-ups at reproduction events. The genealogical structure is obtained by tracing backward-in-time the lines from the alive individuals at a given time to their ancestors. Notice that $\mathcal{P}$ is actually a homogeneous in time Poisson point process (whose intensity depends on the measure Λ), thus the lockdown graph has independent and stationary "increments". To emphasize the role of the measure Λ we will say Λ lockdown graph from now on. Using a sequence of *initial types* $(\xi_0(i))_{i \geq 1}$, i.i.d. $\text{uniform}[0, 1]$, we define the lockdown process $(\xi_t(i), t \in [0, \infty))_{i \geq 1}$ thanks to the Λ lockdown graph $\mathcal{P}$ via the following procedure. For each $i \geq 1$, $t \mapsto \xi_t(i)$ starts from $\xi_0(i)$ and remains constant when the corresponding line does not change of level. At a reproduction event, if level $i \geq 1$ is hit by an arrow, then $\xi_t(i)$ takes the type of the parent, otherwise either there is no push-up at level i and $\xi_t(i)$ remains constant or there is a push-up and $\xi_t(i)$ take the value $\xi_{t-}(j)$, where j is the level (necessarily lower than i) of the line pushed up to level i . The process $(\xi_t(i), t \in [0, \infty))_{i \geq 1}$ will also be designated




Figure 1: A lookdown graph. Each arrow corresponds to a birth event: the level carrying a dot has reproduced on the levels carrying an ending arrow. For example, at time t_1 , level 2 reproduces on levels 5 and 7 while former levels 5, 6 and 7 are pushed up to the next available levels. Note that only finitely many birth events affecting the n first levels occur in any compact interval of time, for each integer n . Furthermore, by tracing back the lineages from a given time, one obtains a Λ coalescent tree.

The corresponding flow of partitions, restricted to $[7]$, would be entirely defined by the partitions $\hat{\Pi}_{t_1-, t_1} = \{\{1\}, \{2, 5, 7\}, \{3\}, \{4\}, \{6\}\}$, $\hat{\Pi}_{t_2-, t_2} = \{\{1, 2, 4\}, \{3\}, \{5\}, \{6\}, \{7\}\}$, $\hat{\Pi}_{t_3-, t_3} = \{\{1\}, \{2, 5\}, \{3\}, \{4\}, \{6\}, \{7\}\}$ and so on.

by $\mathcal{L}_0(\mathcal{P}, (\xi_0(i))_{i \geq 1})$ to emphasize the fact that we started from time 0 with the lookdown graph $\mathcal{P}$ and the initial types $(\xi_0(i))_{i \geq 1}$. Donnelly and Kurtz prove in [12], that the limiting empirical measures

$$\Xi_t(\cdot) := \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \delta_{\xi_t(i)}(\cdot) \text{ for } t \in [0, \infty)$$

exist simultaneously for all $t \geq 0$ almost surely and form a Λ Fleming-Viot process.

Organization of the paper

In Section 2, we recall some notions on Λ coalescents, stochastic flows of bridges and the lookdown representation. In Section 3, we introduce flows of partitions and make the connection with lookdown graphs. In Section 4, we study in detail the Λ Fleming-Viot process, in particular the problem of simultaneous extinction of ancestral types and prove Theorem 1. In addition, we introduce the Eves as an ordering of the ancestral types and give several properties of this sequence. In Section 5, we define the dynamic version of the Eves, that is, the Eves process and, then, we prove Theorem 2. Section 6 is devoted to the proof of Theorem 3. Finally, in Section 7 we reformulate results of Berestycki et al. in [1, 2] on the encoding of the lookdown process associated with the Beta($2 - \alpha, \alpha$) Fleming-Viot into a height process associated with an α -stable branching mechanism, with $\alpha \in (1, 2]$, in terms of the flow of partitions and the Eves process. The upshot of this formulation is that not only the genealogy of the Beta($2 - \alpha, \alpha$) Fleming-Viot process but also its Eves are defined in terms of the height process.

2 Preliminaries

2.1 Coalescent with multiple collisions

Let us recall the definition of the coalescents with multiple collisions, also called Λ coalescents, which are introduced in [17, 18]. As in [6], we denote by $\mathcal{P}_n$ the set of all partitions of $[n] := \{1, 2, \dots, n\}$, with $n \in \mathbb{N} \cup \{\infty\}$. $\mathcal{P}_\infty$ is equipped with the distance $d_{\mathcal{P}}$ defined as follows. For all $\pi, \pi' \in \mathcal{P}_\infty$

$$d_{\mathcal{P}}(\pi, \pi') = 2^{-i} \Leftrightarrow i = \sup\{j \in \mathbb{N} : \pi^{[j]} = \pi'^{[j]}\}$$

where $\pi^{[j]}$ is the restriction of π to $[j]$. The metric space $(\mathcal{P}_\infty, d_{\mathcal{P}})$ is compact.

For each $i \geq 1$, we denote by $\pi(i)$ the i -th block of a given partition $\pi \in \mathcal{P}_\infty$, where the blocks are in the increasing order of their least element. Furthermore, for each $i \geq 1$, we introduce the asymptotic frequency of the i -th block of π as

$$|\pi(i)| := \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \mathbf{1}_{\{j \in \pi(i)\}} \quad (3)$$

when the limit on the r.h.s. exists.

We define the coagulation operator $Coag : \mathcal{P}_\infty \times \mathcal{P}_\infty \rightarrow \mathcal{P}_\infty$ as follows. For any elements $\pi, \pi' \in \mathcal{P}_\infty$, $Coag(\pi, \pi')$ is the partition whose blocks are given by

$$Coag(\pi, \pi')(i) = \bigcup_{j \in \pi'(i)} \pi(j) \quad (4)$$

for every $i \in \mathbb{N}$. This is a Lipschitz-continuous operator and we have

$$Coag(\pi, Coag(\pi', \pi'')) = Coag(Coag(\pi, \pi'), \pi'') \quad (5)$$

for any elements $\pi, \pi', \pi'' \in \mathcal{P}_\infty$, see Section 4.2 in [4] for further details.

Consider a finite measure Λ on $[0, 1]$. A Λ coalescent is a Markov process $(\Pi_t, t \geq 0)$ on $\mathcal{P}_\infty$ started from the partition $0_{[\infty]} := \{\{1\}, \{2\}, \dots\}$ and such that, for each integer $n \geq 2$, its restriction $(\Pi_t^{[n]}, t \geq 0)$ to $\mathcal{P}_n$ is a continuous time Markov chain that evolves by coalescence events whose dynamics is the following. For any integer $2 \leq p \leq n$, consider a partition $\pi \in \mathcal{P}_n$ whose blocks are all singletons except one which has p elements. At rate $\lambda_{n,p}$, $\Pi_t^{[n]}$ jumps to $Coag(\Pi_t^{[n]}, \pi)$ where

$$\lambda_{n,p} = \int_0^1 x^{p-2} (1-x)^{n-p} \Lambda(dx) \quad (6)$$

If $\Pi_t^{[n]}$ has m blocks, the total jump rate of the chain at this time is then

$$\lambda_m = \sum_{p=2}^m \binom{m}{p} \lambda_{m,p} \quad (7)$$

From now on, we will systematically assume that $\Lambda(1) = 0$ to avoid trivial behaviour. Indeed an atom on 1 induces coalescence events involving all the blocks simultaneously. Pitman [17] showed that a Λ coalescent could either come down from infinity (CDI), that is, the process $\#\Pi_t$ is finite at any time

$t > 0$ a.s., or stay infinite, that is, $\#\Pi_t$ is infinite for all $t > 0$ a.s., where $\#\pi$ denotes the number of blocks of a partition π . A necessary and sufficient condition on the measure Λ that ensures the coming down from infinity can be found in [19]. We will denote by $\mathbb{CDI}$ the set of measures Λ for which the Λ coalescent comes down from infinity.

The Λ coalescents obtained with a measure Λ taken as the density of a $\text{Beta}(2 - \alpha, \alpha)$ variable, with $0 < \alpha < 2$, are called $\text{Beta}(2 - \alpha, \alpha)$ coalescents (see [2, 3, 9] for several results about such coalescents). Recall that those densities are given by

$$\Lambda(dx) = \frac{1}{\Gamma(2 - \alpha)\Gamma(\alpha)} x^{1-\alpha}(1 - x)^{\alpha-1} dx \quad (8)$$

Those coalescents come down from infinity iff $\alpha > 1$. The Kingman coalescent is recovered when $\alpha \rightarrow 2$ (we will use the notation $\text{Beta}(0, 2)(dx)$ for $\delta_0(dx)$), whereas the Bolthausen-Sznitman coalescent arises when $\alpha = 1$. Note that the latter is the only $\text{Beta}(2 - \alpha, \alpha)$ coalescent that stays infinite but has no dust (or equivalently, no singleton at any given time $t > 0$) almost surely.

We end this subsection with a result, whose proof is postponed in Section 8, which will be useful in the proof of Theorem 3.

Proposition 2.1 *Let $(\Pi_t, t \geq 0)$ be a Λ coalescent assumed to be càdlàg. Almost surely, for all $t \geq 0$ the partition Π_t admits asymptotic frequencies and for every $i \geq 1$ the process $t \mapsto |\Pi_t(i)|$ is càdlàg.*

2.2 Stochastic flows of bridges

We recall basic definitions and properties of stochastic flows of bridges introduced by Bertoin and Le Gall in [6]. A bridge is a nondecreasing càdlàg process $F = (F(r), r \in [0, 1])$ with values in $[0, 1]$ such that :

- $F(0) = 0, F(1) = 1$
- F has exchangeable increments

Kallenberg [15] shows that for any bridge F , there exists a sequence of nonnegative r.v. $(\beta^i)_{i \in \mathbb{N}}$ with $\beta^1 \geq \beta^2 \geq \dots \geq 0$, and $\sum_{i=1}^{\infty} \beta^i \leq 1$, and a sequence of i.i.d. uniform $[0, 1]$ r.v. $(U^i)_{i \in \mathbb{N}}$ independent of the sequence $(\beta^i)_{i \in \mathbb{N}}$ such that a.s. for every $r \in [0, 1]$,

$$F(r) = \sum_{i=1}^{\infty} \beta^i 1_{\{U^i \leq r\}} + (1 - \sum_{i=1}^{\infty} \beta^i) r \quad (9)$$

From any bridge F and any infinite sequence $(V_p)_{p \geq 1}$ of i.i.d. uniform random variables on $[0, 1]$ independent of F , one can define a random partition $\pi(F, (V_p)_{p \geq 1})$ by

$$i \stackrel{\pi(F, (V_p)_{p \geq 1})}{\sim} j \Leftrightarrow F^{-1}(V_i) = F^{-1}(V_j) \quad (10)$$

In [6] Bertoin and Le Gall define a consistent collection of bridges in order to obtain, using the above construction, a consistent collection of random partitions.

Definition 2.2 A flow of bridges is a collection $(F_{s,t}, -\infty < s \leq t < \infty)$ of bridges such that :

- For every $r < s < t$, $F_{r,t} = F_{s,t} \circ F_{r,s}$ a.s. (cocycle property).
- The law of $F_{s,t}$ only depends on $t - s$. Furthermore, if $s_1 < s_2 < \dots < s_n$ the bridges $F_{s_1,s_2}, F_{s_2,s_3}, \dots, F_{s_{n-1},s_n}$ are independent.
- $F_{0,0} = Id$ and $F_{0,t} \rightarrow Id$ in probability as $t \downarrow 0$, in the sense of Skorohod's topology.

Note that symbol $\circ$ stands for the composition operator of real-valued functions. Given a sequence of i.i.d. uniform $[0, 1]$ variables $(V_p)_{p \geq 1}$, independent of a flow of bridges $(F_{s,t}, -\infty < s \leq t < \infty)$, they prove that the process $(\pi(F_{-t,0}, (V_p)_{p \geq 1}))_{t \geq 0}$ is an exchangeable coalescent, see Theorem 1 in [6]. In the particular case of a Λ coalescent, $(F_{s,t}, -\infty < s \leq t < \infty)$ is called a Λ flow of bridges.

Remark that the processes $(F_{0,t}, t \geq 0)$ and $(F_{-t,0}, t \geq 0)$ do not have the same distribution, even if for all $t \geq 0$

$$F_{0,t} \stackrel{(d)}{=} F_{-t,0} \quad (11)$$

Roughly speaking, the process $(F_{-t,0}, t \geq 0)$ provides the genealogy of the population, while the process $(F_{0,t}, t \geq 0)$ describes the evolution forward-in-time. From the latter, Bertoin and Le Gall define the Λ Fleming-Viot process as follows. Denote by $\mathcal{M}_1$ the space of all probability measures on $[0, 1]$, equipped with its weak topology. The Λ Fleming-Viot process is defined as the $\mathcal{M}_1$ -valued process $(\rho_{0,t}, t \in [0, \infty))$ where

$$\rho_{0,t}([0, x]) = F_{0,t}(x), \text{ for all } x \in [0, 1]$$

Bertoin and Le Gall in [6] prove that this process is a Markov process with a Feller semigroup which is characterized by a martingale problem (based on a duality argument with the Λ coalescent) that we do not recall here. Note that, from the stationarity property verified by a flow of bridges, for every $s \in \mathbb{R}$ the process $(\rho_{s,t}, t \in [s, \infty))$ is also a Λ Fleming-Viot process, where the probability measures $\rho_{s,t}$ are defined from the bridge $F_{s,t}$ as above.

Since the semigroup is Feller, each process $(\rho_{s,t}, t \in [s, \infty))$ admits a càdlàg modification still denoted $(\rho_{s,t}, t \in [s, \infty))$ for simplicity. The collection of bridges associated to these càdlàg modifications are also still denoted $(F_{s,t}, s \leq t)$.

Remark 2.3 One should realize that a stochastic flow of bridges F , except when it arises from a Poissonian construction with elementary bridges (see Section 5 in [6]), may have many irregularities : the cocycle property does not necessarily hold simultaneously for all triplets $r < s < t$, almost surely. This will be a difficulty throughout this work : all the objects defined pathwise from the flow will suffer from those irregularities and will need themselves to be regularized, as we will see later for the flow of partitions.

2.3 The lockdown process

Let us recall the definition of the lockdown process, introduced by Donnelly and Kurtz in [11, 12] and generalized to the case of Ξ -coalescents in [10]. As explained in the introduction, this process relies on

the so-called lookdown graph; we refer to the introduction for a heuristic definition of this object, here we will give a precise definition. Let us first define some notation. For each $n \in \{2, 3, \dots, \infty\}$, let $\mathcal{S}_n$ be the subset of $\{0, 1\}^n$ whose elements have at least two coordinates $1 \leq i < j \leq n$ equal to 1. For all integer $n \geq 2$, we denote by $[u]_n := (u^1, \dots, u^n)$ the restriction of an element $u = (u^1, \dots, u^n, u^{n+1}, \dots) \in \mathcal{S}_\infty$ to its n first coordinates. Remark that $[u]_n$ is not necessarily an element of $\mathcal{S}_n$. Thus, for a given subset A of $\mathcal{S}_\infty$, we set

$$A|_{\mathcal{S}_n} := \{[u]_n : u \in A\} \cap \mathcal{S}_n$$

Note that $A|_{\mathcal{S}_n}$ can be empty.

Definition 2.4 *A deterministic lookdown graph is a deterministic point collection p on $\mathbb{R} \times \mathcal{S}_\infty$ such that for each $n \in \mathbb{N}$, for all $s \leq t$, $p|_{[s,t] \times \mathcal{S}_n}$ has finitely many points.*

The point collection p should be seen as a collection of points $(t, u) \in \mathbb{R} \times \mathcal{S}_\infty$ called reproduction events, where t designates the reproduction time and u determines the individuals that participate to this event. More precisely, the set

$$I_{t,u} := \{i \geq 1 : u^i = 1\} \quad (12)$$

is called the set of individuals that participate to the reproduction event.

Fix $s \in \mathbb{R}$ and consider a vector of initial types $(\xi_{s,s}(i))_{i \geq 1} \in [0, 1]^\mathbb{N}$. We define a particle system $\xi_{s,t} = (\xi_{s,t}(1), \xi_{s,t}(2), \dots), t \in [s, \infty)$ with values in $[0, 1]^\infty$, as follows :

- The initial values are given by $(\xi_{s,s}(i))_{i \geq 1}$.
- At any birth event $(t, u) \in p$ with $t > s$, for each $i \geq 1$ the function $r \mapsto \xi_{s,r}(i)$ evolves as follows

$$\begin{cases} \xi_{s,t}(i) = \xi_{s,t-(\min(I_{t,u}))} & \text{for all } i \in I_{t,u} \\ \xi_{s,t}(i) = \xi_{s,t-(i - (\#\{I_{t,u} \cap [i]\} - 1) \vee 0)} & \text{for all } i \notin I_{t,u} \end{cases} \quad (13)$$

It is called a deterministic lookdown function. Remark that this is the deterministic lookdown construction with push-up, that is, instead of killing particles located at birth levels, they are pushed up to the next available level (see Figure 1). It corresponds to the modified lookdown construction of Donnelly and Kurtz [12]. We thus introduce the following notation.

Definition 2.5 *We denote by $\mathcal{L}_s(p, (\xi_{s,s}(i))_{i \geq 1})$ the deterministic lookdown function $(\xi_{s,t}(i), t \in [s, \infty))_{i \geq 1}$ started at time s with the point collection p and the initial types $(\xi_{s,s}(i))_{i \geq 1}$.*

When it exists, for every $t \in [s, \infty)$ let

$$\Xi_{s,t}(\cdot) := \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \delta_{\xi_{s,t}(i)}(\cdot) \quad (14)$$

be the limiting empirical measure of this deterministic lookdown function taken at time t .

Definition 2.6 *We denote by $\mathcal{E}_s(p, (\xi_{s,s}(i))_{i \geq 1})$ the collection of limiting empirical measures $(\Xi_{s,t}, t \in [s, \infty))$ started at time s with the point collection p and the initial types $(\xi_{s,s}(i))_{i \geq 1}$, when it exists.*

We now explain how one can define a random lockdown process such that its limiting empirical measures are almost surely defined, and form a Λ Fleming-Viot process, where Λ is a finite measure on $[0, 1]$. We take $\mathcal{P}$ as a Poisson point process on $\mathbb{R} \times \mathcal{S}_\infty$ with intensity measure $dt \otimes (\mu_K + \mu_\Lambda)$ where μ_K and μ_Λ are defined as follows.

The intensity measure μ_Λ on $\mathcal{S}_\infty$ corresponds to resampling events with positive frequency, that is, reproduction events involving a positive proportion of individuals. Let $\nu^x(\cdot)$ be the distribution on $\mathcal{S}_\infty$ of a sequence of i.i.d. Bernoulli random variables with parameter x , for each $x \in (0, 1)$.

$$\mu_\Lambda(\cdot) := \int_{(0,1)} x^{-2} \Lambda(dx) \nu^x(\cdot) \quad (15)$$

We now define the intensity measure for Kingman's reproduction events, that is, reproduction events involving only two individuals at once. For each $1 \leq i < j$, let $s_{i,j}$ be the element of $\mathcal{S}_\infty$ that has only two coordinates equal to 1: i and j . We define the measure μ_K on $\mathcal{S}_\infty$ by

$$\mu_K(\cdot) := \Lambda(0) \sum_{1 \leq i < j} \delta_{s_{i,j}}(\cdot) \quad (16)$$

Definition 2.7 *A lockdown graph associated with the measure Λ - or Λ lockdown graph in short - is a Poisson point process $\mathcal{P}$ on $\mathbb{R} \times \mathcal{S}_\infty$ with intensity measure $dt \otimes (\mu_K + \mu_\Lambda)$.*

Remark 2.8 *Consider such a Poisson point process. Almost surely, for all $s < t$ and every $n \in \mathbb{N}$, $\mathcal{P}|_{[s,t] \times \mathcal{S}_n}$ has finitely many points.*

Consider a lockdown graph $\mathcal{P}$ associated with the measure Λ . Fix a time $s \in \mathbb{R}$ and a sequence of i.i.d. uniform $[0, 1]$ r.v. $(\xi_{s,s}(i))_{i \geq 1}$. Donnelly and Kurtz prove in Lemma 3.5 of [12] that $\mathcal{L}_s(\mathcal{P}, (\xi_{s,s}(i))_{i \geq 1})$ admits a limiting empirical measure $\Xi_{s,t}$ simultaneously for all $t \in [s, \infty)$, almost surely. Moreover, the process $(\Xi_{s,t}, t \in [s, \infty))$ is a càdlàg Λ Fleming-Viot process.

The lockdown graph can then be used to define a collection of lockdown processes indexed by their starting time $s \in \mathbb{R}$, if we are given for each of them a sequence of initial types. Hence their genealogies are coupled, but their types are not, unless the sequences of initial types were suitably coupled as we will see later. In particular, to obtain a flow of bridges, the coupling of the initial types is not straightforward.

We end this subsection with a definition that will be useful in the sequel. Consider a lockdown process $(\xi_t(i), t \in [0, \infty))_{i \geq 1} = \mathcal{L}_0(\mathcal{P}, (\xi_0(i))_{i \geq 1})$ (for simplicity, we write ξ_t instead of $\xi_{0,t}$ to alleviate notation).

Definition 2.9 *We define $Y_t(i)$ as the lowest level at time t that carries the type $\xi_0(i)$.*

$$Y_t(i) := \inf\{j \geq 1 : \xi_t(j) = \xi_0(i)\} \quad (17)$$

3 Flows of partitions

In this section, we introduce flows of partitions. We will see that this new object will provide an efficient and elegant way to encode lockdown graphs. In addition, it possesses both aspects of the two representations we intend to couple : as a flow it verifies a cocycle property and has independent and stationary increments, while the use of partitions allows to restrict to a finite number of individuals sampled into the current population (as does the lockdown representation).

We begin by giving a deterministic definition of a flow of partitions and we show that, under a technical assumption, it is equivalent with a deterministic lockdown graph. Then, we randomize these objects and introduce stochastic flows of partitions. Many results are technical so their proofs are postponed to Section 8. On first reading one can restrict oneself to Subsection 3.1 and Definition 3.6.

3.1 Deterministic flows of partitions

Definition 3.1 *A deterministic flow of partitions is a collection $(\hat{\pi}_{s,t}, -\infty < s \leq t < \infty)$ of partitions such that*

- *For every $r < s < t$, $\hat{\pi}_{r,t} = \text{Coag}(\hat{\pi}_{s,t}, \hat{\pi}_{r,s})$ (cocycle property).*
- *For every $s \in \mathbb{R}$, $\lim_{r \uparrow s} \lim_{t \uparrow s} \hat{\pi}_{r,t} =: \hat{\pi}_{s-,s-} = 0_{[\infty]}$ (left regularity).*
- *For every $s \in \mathbb{R}$, $\lim_{t \downarrow s} \hat{\pi}_{s,t} = \hat{\pi}_{s,s} = 0_{[\infty]}$ (right regularity).*

Furthermore, if for all $s \in \mathbb{R}$, $\hat{\pi}_{s-,s}$ has at most one unique non-singleton block, then we say that $\hat{\pi}$ is a deterministic flow of partitions without simultaneous mergers.

We begin with an elementary lemma. We use the notation $\pi^{[n]}$ to designate the restriction to $[n]$ of a given partition $\pi \in \mathcal{P}_\infty$.

Lemma 3.2 *For all $s \in \mathbb{R}$ and $n \in \mathbb{N}$, there exists $\epsilon > 0$ such that*

$$\begin{aligned} \hat{\pi}_{r,t}^{[n]} &= 0_{[n]} \text{ if } s \leq r \leq t < s + \epsilon \\ \hat{\pi}_{r,t}^{[n]} &= 0_{[n]} \text{ if } s - \epsilon < r \leq t < s \end{aligned}$$

Proof This is a consequence of three facts: the left and right regularities of a deterministic flow of partitions, the cocycle property and the finiteness of the set $\mathcal{P}_n$. □

As a consequence, one can easily deduce that $\hat{\pi}_{s-,s}$ is well defined for each $s \in \mathbb{R}$ (and thus, the last sentence of the preceding definition makes sense). In addition we have the following corollary, whose proof can be deduced from the preceding lemma.

Corollary 3.3 *For all $s \in \mathbb{R}$, $(\hat{\pi}_{s,t}, t \in [s, \infty))$ is a càdlàg $\mathcal{P}_\infty$ -valued function and $(\hat{\pi}_{s-r,s}, r \in [0, \infty))$ is a ladcàg $\mathcal{P}_\infty$ -valued function.*

Such flows of partitions are actually related with deterministic lockdown graphs, let us show how (see also Figure 1). We introduce, for each $n \in \mathbb{N} \cup \{\infty\}$, $\mathcal{P}_n^*$ as the subset of $\mathcal{P}_n$ whose elements have a unique non-singleton block. The key tool is the map $g_n : \mathcal{S}_n \rightarrow \mathcal{P}_n^*$ that translates a reproduction event in terms of partitions as follows. For a reproduction event $u = \{u^1, \dots, u^n\} \in \mathcal{S}_n$, we denote by $I := \{i \in [n] : u^i = 1\}$ the corresponding set of individuals that participate. The partition $g_n(u)$ is then defined as the element of $\mathcal{P}_n^*$ whose unique non-singleton block is I , that is, for all $i, j \in [n]$

$$i \stackrel{g_n(u)}{\sim} j \Leftrightarrow u^i = u^j = 1 \quad (18)$$

Plainly, g_n is a bijection from $\mathcal{S}_n$ to $\mathcal{P}_n^*$.

Proposition 3.4 *There exists a one-to-one correspondence between the set of deterministic flows of partitions without simultaneous mergers and the set of deterministic lockdown graphs.*

Proof Consider a deterministic lockdown graph p . For each $n \in \mathbb{N}$ and every $s < t$, let $(t_m, u_m)_{1 \leq m \leq q}$ denote the finitely many atoms of $p|_{(s,t] \times \mathcal{S}_n}$ in the increasing order of their time coordinate and set

$$\hat{\pi}_{s,t}^{[n]} := \text{Coag}(g_n(u_q), \text{Coag}(g_n(u_{q-1}), \dots, \text{Coag}(g_n(u_2), g_n(u_1)) \dots)) \quad (19)$$

and $\hat{\pi}_{s,s}^{[n]} := 0_{[n]}$. Obviously, the collection of partitions $(\hat{\pi}_{s,t}^{[n]}, n \in \mathbb{N})$ is compatible and defines by a projective limit a unique partition $\hat{\pi}_{s,t}$ such that its restriction to $[n]$ is $\hat{\pi}_{s,t}^{[n]}$ for each $n \in \mathbb{N}$. Thus, it is straightforward to verify that the collection of partitions $(\hat{\pi}_{s,t}, -\infty < s \leq t < \infty)$ is a deterministic flow of partitions without simultaneous mergers.

Conversely consider a deterministic flow of partitions $\hat{\pi}$ without simultaneous mergers. We define the collection of its jumps $p := \bigcup_{s: \pi_{s-,s} \neq 0_{[\infty]}} \{s, g_\infty^{-1}(\hat{\pi}_{s-,s})\}$ which is a point collection on $\mathbb{R} \times \mathcal{S}_\infty$. We have to check that the restriction to $\mathbb{R} \times \mathcal{S}_n$ of this point collection does not accumulate near any point in $\mathbb{R}$. This follows from Lemma 3.2. Therefore, we conclude that p is a deterministic lockdown graph.

Finally, it is easy to show that the composition of these two procedures is the identity. This entails the one-to-one correspondence. $\square$

The interest of this correspondence is that the flow of partitions entirely encodes the genealogical relationships of the lockdown graph. Indeed, consider a deterministic lockdown graph p and let $\hat{\pi}$ be the deterministic flow of partitions associated via the preceding bijection. Set $(\xi_{0,t}(i), t \in [0, \infty)) = \mathcal{L}_0(p, (\xi_{0,0}(i))_{i \geq 1})$, for a given sequence of initial types $(\xi_{0,0}(i))_{i \geq 1}$.

Lemma 3.5 *For all $t \in [0, \infty)$ and all $i, j \in \mathbb{N}$*

$$\xi_{0,t}(j) = \xi_{0,0}(i) \Leftrightarrow j \in \hat{\pi}_{0,t}(i) \quad (20)$$

Proof This is a simple consequence of the properties of the coagulation operator and of the definition of the lockdown process. $\square$

In words, this means that each block $\hat{\pi}_{0,t}(i)$ is the set of individuals alive at time t whose ancestor at time 0 had type $\xi_{0,0}(i)$, for every $i \geq 1$. This lemma allows us to introduce the notation $\mathcal{L}_0(\hat{\pi}, (\xi_{0,0}(i))_{i \geq 1})$ which defines the same object as $\mathcal{L}_0(p, (\xi_{0,0}(i))_{i \geq 1})$.

3.2 Stochastic flows of partitions

In this subsection, we introduce stochastic flows of partitions which are new objects of independent interest. We present the construction from a lookdown graph of a stochastic flow of partitions, then we show how a given stochastic flow of partitions encodes pathwise a lookdown graph.

Definition 3.6 *A stochastic flow of partitions is a collection of random partitions $\hat{\Pi} = (\hat{\Pi}_{s,t}, -\infty < s \leq t < \infty)$ that enjoys the following properties:*

- For every $r < s < t$, $\hat{\Pi}_{r,t} = Coag(\hat{\Pi}_{s,t}, \hat{\Pi}_{r,s})$ a.s. (cocycle property).
- $\hat{\Pi}_{s,t}$ is an exchangeable random partition whose law only depends on $t - s$. Furthermore, for any $s_1 < s_2 < \dots < s_n$ the partitions $\hat{\Pi}_{s_1, s_2}, \hat{\Pi}_{s_2, s_3}, \dots, \hat{\Pi}_{s_{n-1}, s_n}$ are independent.
- $\hat{\Pi}_{0,0} = 0_{[\infty]}$ and $\hat{\Pi}_{0,t} \rightarrow 0_{[\infty]}$ in probability as $t \downarrow 0$, for the distance $d_{\mathcal{P}}$.

Moreover, when the process $(\hat{\Pi}_{-t,0}, t \geq 0)$ is a Λ coalescent, we say that $\hat{\Pi}$ is associated with the measure Λ or, in short, is a Λ flow of partitions.

Remark 3.7 *It is worth noting that flows of partitions have been introduced separately by Foucart [14] as a population model for Ξ Fleming-Viot processes with immigration.*

Note that the processes $(\hat{\Pi}_{0,t}, 0 \leq t < \infty)$ and $(\hat{\Pi}_{-t,0}, 0 \leq t < \infty)$ do not have the same distribution, even if their one-dimensional marginals are the same. Introduce the following filtration.

$$\mathcal{F}_t^{\hat{\Pi}} := \sigma\{\hat{\Pi}_{r,s}, r \leq s \leq t\}$$

Lemma 3.8 *The process $(\hat{\Pi}_{0,t}, 0 \leq t < \infty)$ is a $\mathcal{F}^{\hat{\Pi}}$ Markov process taking values in $\mathcal{P}_{\infty}$ with a Feller semigroup.*

Proof Consider $(\hat{\Pi}_{0,t}, 0 \leq t < \infty)$. The very definition of stochastic flows of partitions ensures that this process is Markov with a semigroup Q_t such that for every $\pi \in \mathcal{P}_{\infty}$

$$Q_t(\pi, \cdot) \stackrel{(d)}{=} Coag(\hat{\Pi}_{0,t}, \pi) \tag{21}$$

Clearly $Q_t \circ Q_s = Q_{t+s}$. Recall that $(\mathcal{P}_{\infty}, d_{\mathcal{P}})$ is compact and consider a bounded continuous function $f : \mathcal{P}_{\infty} \rightarrow \mathbb{R}$.

$$Q_t f(\pi) = \mathbb{E}[f(Coag(\hat{\Pi}_{0,t}, \pi))]$$

Since $Coag$ is continuous, by dominated convergence we get that $Q_t f$ is a bounded continuous function. Notice that $\hat{\Pi}_{0,t} \rightarrow 0_{[\infty]}$ in probability. Then, for any $\pi \in \mathcal{P}_{\infty}$

$$Q_t f(\pi) = \mathbb{E}[f(Coag(\hat{\Pi}_{0,t}, \pi))] \xrightarrow[t \downarrow 0]{} f(\pi)$$

This implies the Feller property of Q . □

One should pay attention to the fact that the trajectories of a given stochastic flow of partitions $\hat{\Pi}$ are not necessarily deterministic flows of partitions in the sense of our definition. Indeed, the cocycle property does not necessarily hold simultaneously for every triplet $r < s < t$ almost surely (compare with Remark 2.3). Hence the trajectories are not necessarily deterministic flows of partitions, and so, the one-to-one correspondence of the previous subsection cannot be directly applied to obtain deterministic lockdown graphs; but this difficulty will be overcome thanks to the regularization procedure (see below). Conversely, starting from a Λ lockdown graph, the one-to-one correspondence can be applied as follows.

Construction from a lockdown graph. Fix a finite measure Λ on $[0, 1]$ and consider a Λ lockdown graph $\mathcal{P}$, that is, a Poisson point process on $\mathbb{R} \times \mathcal{S}_\infty$ with intensity $dt \otimes (\mu_K + \mu_\Lambda)$ (recall the definitions of Subsection 2.3). Note that for $\mathbb{P}$ -a.a. $\omega \in \Omega$, the point collection $\mathcal{P}(\omega)|_{[s,t] \times \mathcal{S}_n}$ has finitely many points for all $s < t$ and all $n \geq 1$. So without loss of generality, we can assume that it holds for all $\omega \in \Omega$. Therefore, one can define a deterministic flow of partitions $\hat{\Pi}^\mathcal{P}(\omega)$ using Proposition 3.4 and the point collection $\mathcal{P}(\omega)$ for each $\omega \in \Omega$.

Proposition 3.9 *The process $\hat{\Pi}^\mathcal{P}$ is a Λ flow of partitions.*

Proof Fix $r \leq s \leq t$. For all $n \in \mathbb{N}$, we easily deduce from the definition that

$$\hat{\Pi}_{r,t}^{\mathcal{P},[n]} = Coag(\hat{\Pi}_{s,t}^{\mathcal{P},[n]}, \hat{\Pi}_{r,s}^{\mathcal{P},[n]})$$

This implies that $\hat{\Pi}_{r,t}^\mathcal{P} = Coag(\hat{\Pi}_{s,t}^\mathcal{P}, \hat{\Pi}_{r,s}^\mathcal{P})$.

The independence and continuity properties are straightforward from the definition. Finally the Poissonian construction of coalescent processes (see [4]) ensures that $(\hat{\Pi}_{-t,0}^\mathcal{P}, t \geq 0)$ is a Λ coalescent. $\square$

Remark 3.10 *Remark that the trajectories of a stochastic flow of partitions constructed from a lockdown graph are deterministic flows of partitions.*

This concludes the Construction from a lockdown graph.

We now consider a Λ flow of partitions. The goal is to define an associated Λ lockdown graph using the one-to-one correspondence. But as we said above, the trajectories of the flow of partitions are not necessarily deterministic flows of partitions. Therefore it is necessary to define a modification of this flow, such that the trajectories of this modification are deterministic flows of partitions a.s. The reason that motivates this technical discussion is that we will identify in Section 5 a stochastic flow of partitions embedded into a flow of bridges. Then to use the correspondence with lockdown graphs, we will need to work with trajectories that are deterministic flows of partitions.

Regularization of a stochastic flow of partitions. Consider a Λ flow of partitions $\hat{\Pi}$. The strategy is to restrict to the rational marginals of the flow of partitions, and to prove that they verify the properties of a deterministic flow of partitions outside an event of zero probability. Then, we take right and left limits on this object and prove that we recover a modification of the initial flow whose trajectories are deterministic flows of partitions almost surely. The proofs are postponed to Section 8.

Lemma 3.11 Consider the restriction $(\hat{\Pi}_{s,t}, s, t \in \mathbb{Q})$ of the flow of partitions to its rational marginals. Then there exists an event $\Omega_{\hat{\Pi}}$ of probability 1 such that on this event, the trajectories are deterministic flows of partitions (restricted to the rational marginals). More precisely, on $\Omega_{\hat{\Pi}}$ the cocycle property is verified for all rational marginals and for every $s \in \mathbb{R}$ and $n \in \mathbb{N}$, there exists $\epsilon(\omega) > 0$ such that

$$\begin{aligned}\hat{\Pi}_{p,q}^{[n]} &= 0_{[n]} \text{ if } p, q \in [s, s + \epsilon) \cap \mathbb{Q} \\ \hat{\Pi}_{p,q}^{[n]} &= 0_{[n]} \text{ if } p, q \in (s - \epsilon, s) \cap \mathbb{Q}\end{aligned}$$

We now define for every $s \leq t \in \mathbb{R}$ the partition $\tilde{\Pi}_{s,t}$ on the event $\Omega_{\hat{\Pi}}$ as follows.

Lemma 3.12 On the event $\Omega_{\hat{\Pi}}$, the following random partition is well-defined.

$$\tilde{\Pi}_{s,t} := \begin{cases} \lim_{v \downarrow t, v \in \mathbb{Q}} \hat{\Pi}_{s,v} & \text{if } s \in \mathbb{Q} \\ \lim_{r \downarrow s, r \in \mathbb{Q}} \hat{\Pi}_{r,t} & \text{if } t \in \mathbb{Q} \\ \text{Coag}(\tilde{\Pi}_{q,t}, \tilde{\Pi}_{s,q}) & \text{for any arbitrary rational } q \in (s, t) \text{ if } s, t \notin \mathbb{Q} \\ 0_{[\infty]} & \text{if } t = s \end{cases} \quad (22)$$

Furthermore, for every $r < s < t$, $\tilde{\Pi}_{r,t} = \text{Coag}(\tilde{\Pi}_{s,t}, \tilde{\Pi}_{r,s})$.

On the complementary of $\Omega_{\hat{\Pi}}$, set any arbitrary value to $\tilde{\Pi}_{s,t}$.

Proposition 3.13 The collection of partitions $\tilde{\Pi}$ is a modification of $\hat{\Pi}$, that is, for every $s \leq t$, a.s. $\tilde{\Pi}_{s,t} = \hat{\Pi}_{s,t}$. Furthermore, for each $\omega \in \Omega_{\hat{\Pi}}$, $\tilde{\Pi}(\omega)$ is a deterministic flow of partitions.

For each $\omega \in \Omega_{\hat{\Pi}}$, let $\mathcal{P}(\omega)$ be the deterministic lookdown graph obtained from $\hat{\Pi}(\omega)$ by applying Proposition 3.4. On the complementary, set any arbitrary values to $\mathcal{P}$.

Proposition 3.14 The point collection $\mathcal{P}$ is a Λ lookdown graph.

Proof The distribution of this point process is obtained from the definition of the stochastic flow of partitions. $\square$

Remark 3.15 From a stochastic flow of partitions, we have been able to define a regularized modification. Note that this operation does not seem possible for a stochastic flow of bridges. Indeed, a key argument in our proof relies on the continuity of the coagulation operator whereas this property does not hold with the composition operator for bridges.

4 Lambda Fleming-Viot and its ancestral types

4.1 Simultaneous extinction of ancestral types

In this section, we study some properties of Λ Fleming-Viot processes. We denote by $(\rho_t, t \geq 0)$ such a process assumed to be càdlàg (this Markov process enjoys the Feller property), started from the uniform distribution on $[0, 1]$. A point $x \in [0, 1]$ is called *ancestral type* if there exists $t > 0$ such that

$\rho_t(\{x\}) > 0$. We stress that the collection of ancestral types is at most countable. Indeed, in the look-down representation, there is a countable number of initial types. Since the atomic support of the Λ Fleming-Viot process is included into the set of the initial types of its lookdown representation, the claim follows.

When ρ does not charge an ancestral type anymore from a given time d , we say that this ancestral type becomes extinct at d . If all ancestral types but one are extinct, we say that the remaining ancestral type has fixed.

The process of asymptotic frequencies of an ancestral type has two possible behaviours: either it leaves continuously 0 at time 0+, or it leaves 0 on a positive jump at a positive time $t > 0$. Therefore we say that an ancestral type $x \in [0, 1]$ *leaves continuously* 0 if $\rho_\epsilon(\{x\}) > 0$ for all $\epsilon > 0$ small enough, and we say that it *emerges from dust* at time $t > 0$ if

$$\forall s \in [0, t), \rho_s(x) = 0, \rho_t(x) > 0 \quad (23)$$

The following proposition is a compilation of results; part of them are a consequence of known facts (see [6]).

Proposition 4.1 *If $\Lambda \in \mathbb{CDI}$, then almost surely :*

- *The measure ρ_t has no continuous part, for all $t > 0$.*
- *Only a finite number of ancestral types have not become extinct at any time $t > 0$.*
- *The primitive Eve fixes in finite time.*

If $\Lambda \notin \mathbb{CDI}$, none of the ancestral types become extinct in finite time, almost surely. Moreover, the following dichotomy occurs :

- *The Λ coalescent has no dust. Then all the ancestral types leave continuously 0.*
- *The Λ coalescent has dust. Then each ancestral type emerges from dust at a positive time.*

Proof If $\Lambda \in \mathbb{CDI}$, then a Λ coalescent $(\Pi_t, t \geq 0)$ has no singleton almost surely [17]. From Equation (11), it is easy to deduce that for all $t > 0$, ρ_t has no continuous part and

$$\#\rho_t \stackrel{(d)}{=} \#\Pi_t \quad (24)$$

where $\#\rho_t$ is the number of atoms of the measure ρ_t , while $\#\Pi_t$ is the number of blocks of the partition Π_t . Since $\#\Pi_t$ reaches 1 in finite time a.s., we deduce that the primitive Eve fixes in finite time. Indeed, $\{\#\rho_t = 1\}_{t>0}$ is a nested collection of events, such that

$$\mathbb{P}(\{\#\rho_t = 1\}) = \mathbb{P}(\{\#\Pi_t = 1\}) \xrightarrow{t \rightarrow \infty} 1$$

Suppose $\Lambda \notin \mathbb{CDI}$. Let us use the lookdown representation of the Λ Fleming-Viot process. Each ancestral type is carried by a certain level i at time 0. Denote by $Y_t(i)$ the lowest level at time t that carries type $\xi_0(i)$ (see Definition 2.9). We claim that $(Y_t(i), t \geq 0)$ does not reach ∞ in finite time. Indeed, the contrary would imply that, with positive probability, only types $\xi_0(1), \xi_0(2), \dots, \xi_0(i-1)$ have not become extinct at a certain finite time and we would deduce that the number of blocks of the Λ coalescent would have a positive probability to be finite at this time, which contradicts our assumption. Hence, none of the ancestral types become extinct.

Finally, let us prove the last dichotomy. Fix $i \geq 1$, and consider the process $(Y_t(i), t \geq 0)$. The presence of dust in the coalescent is equivalent with the fact that the measure $x^{-1}\Lambda(dx)$ is finite [17]. Then, if the coalescent has dust, the first reproduction event involving the line Y as a parent arrives at a strictly positive time and then, this ancestral type emerges from dust. Otherwise, there is no first reproduction event involving the line Y as a parent but an accumulation of infinitely many small (in terms of frequency) such reproduction events. Then its asymptotic frequency leaves 0 continuously at time $0+$. $\square$

We now focus on the coming down from infinity case, and consider the following event

$$E := \{\text{There exists } t > 0 \text{ s.t. two ancestral types become extinct simultaneously at time } t\} \quad (25)$$

The objective of what follows is to prove Theorem 1. This is done thanks to Lemma 4.2 and Proposition 4.4.

Lemma 4.2 *When $\Lambda \in \mathbb{CDI}$, the event E is trivial, that is, $\mathbb{P}(E) \in \{0, 1\}$.*

Remark 4.3 *If $\Lambda \notin \mathbb{CDI}$, the lemma still holds and the event E has probability 0 since none of the ancestral types get extinct.*

Proof Consider a Λ flow of partitions $\hat{\Pi}$ with trajectories assumed to be deterministic flows of partitions (this can be obtained via the Poissonian construction of Subsection 3.2). Let $(\xi_0(i))_{i \geq 1}$ be an independent sequence of i.i.d. uniform $[0, 1]$ r.v. Set $(\xi_t(i), t \geq 0)_{i \geq 1} := \mathcal{L}_0(\hat{\Pi}, (\xi_0(i))_{i \geq 1})$ and $(\rho_t, t \geq 0) := \mathcal{E}_0(\hat{\Pi}, (\xi_0(i))_{i \geq 1})$. We know that ρ is a Λ Fleming-Viot process. We stress that E is independent of $(\xi_0(i))_{i \geq 1}$ and only depends on the flow of partitions. Thus, introduce the filtration $\mathcal{F}$ as follows.

$$\mathcal{F}_t := \sigma\{\hat{\Pi}_{r,s}; 0 \leq r \leq s \leq t\} \text{ for all } t \geq 0$$

One easily remarks that $\mathcal{F}_{0+}$ is a trivial σ -field under $\mathbb{P}$. Set $d^i := \inf\{t \geq 0 : Y_t(i) = \infty\}$, that is, the death time of the i -th initial type in the lookdown representation (see Definition 2.9), which is a stopping time of the filtration $\mathcal{F}$. Since $d^i \downarrow 0$ almost surely as $i \rightarrow \infty$, we deduce that $\bigcap_{i \geq 1} \mathcal{F}_{d^i} = \mathcal{F}_{0+}$. For each $i \geq 1$, define the following event

$$E_i := \{\text{There exists } t \leq d^i \text{ s.t. two ancestral types become extinct simultaneously at time } t\} \quad (26)$$

and $E_\infty := \bigcap_{i \geq 1} E_i$. Clearly, $E_\infty \in \mathcal{F}_{0+}$ so it has probability 0 or 1 under $\mathbb{P}$.

Case 1 : $\mathbb{P}(E_\infty) = 1$ Since $E_\infty \subset E$, we deduce that $\mathbb{P}(E) = 1$.

Case 2 : $\mathbb{P}(E_\infty) = 0$ Suppose there exists $n \geq 1$ such that $\mathbb{P}(E_n) > 0$. It implies that there exists $i \geq n$ and $p > 0$ such that

$$\mathbb{P}(\{d^i = d^{i+1}\}) = p \quad (27)$$

For each $k \geq i$, let

$$\tau_k := \inf\{t \geq 0 : Y_t(i) \geq k\} \quad (28)$$

which is a stopping-time of the filtration $\mathcal{F}$. Remark that

$$\{d^i = d^{i+1}\} = \{Y(i) \text{ and } Y(i+1) \text{ reach } \infty \text{ simultaneously}\}$$

By applying the Markov property at time τ_k (and the fact that the distribution of the lookdown graph is invariant by shift in time), we deduce that

$$\mathbb{P}(\{d^i = d^{i+1}\}) = \mathbb{P}(\{d^{Y_{\tau_k}(i)} = d^{Y_{\tau_k}(i+1)}\}) \quad (29)$$

$$\leq \mathbb{P}(E_k) \quad (30)$$

Hence, for each $k \geq i$, $\mathbb{P}(E_k) \geq p$. Taking the limit when $k \uparrow \infty$, we deduce that $\mathbb{P}(E_\infty) \geq p$, which contradicts our assumption. This implies that for each $i \geq 1$

$$\mathbb{P}(\{d^i = d^{i+1}\}) = 0 \quad (31)$$

which in turn implies that $\mathbb{P}(E) = 0$.

This ends the proof. □

We now determine the probability of E for some important measures Λ .

Proposition 4.4 *For a Beta(2 - α , α) Fleming-Viot process, with $1 < \alpha \leq 2$, we have $\mathbb{P}(E) = 0$.*

In Section 7, we recall the encoding of the lookdown process into a height process associated with an α -stable branching mechanism (see [2, 9, 13]). We use the notation introduced in that section for the proof of the proposition.

Proof Consider all the excursions of H above level 0. These excursions are distributed according to a Poisson point process on $[0, Z_0 = 1] \times \mathcal{E}_+$ where $\mathcal{E}_+$ stands for the set of positive excursions, with intensity measure $dt \otimes \nu^{exc}$ where ν^{exc} is the excursion measure of the height process. Since the measure ν^{exc} gives a finite mass to the set of positive continuous functions whose supremum is greater than any given threshold $\epsilon > 0$, one can order those excursions by decreasing height. The lookdown representation of [2] can then be restated as follows. Consider a sequence of i.i.d. uniform $[0, 1]$ random variables $(\xi_0(i), i \geq 1)$, and associate each type $\xi_0(i)$ to the i -th highest excursion of H above level 0. Then at any time $t > 0$, $\xi_t(j)$ is defined as the type of the j -th highest excursion above level $U(t)$, for each $j \geq 1$, where the type of the j -th highest excursion above level $U(t)$ is the type of the (unique) excursion above 0 in which it is embedded. Results of [2] ensure that $(\xi_t(i), t \geq 0)_{i \geq 1}$ is a lookdown representation of a Beta(2 - α , α) Fleming-Viot process.

It is then sufficient to prove that two distinct particles $(Y_t(i), t \geq 0)$ and $(Y_t(j), t \geq 0)$, with $1 < i < j$, never reach ∞ simultaneously (see Definition 2.9) in order to prove our proposition. This is equivalent to saying that the i -th and j -th highest excursions above level 0 do not die simultaneously in the coalescent time scale. But it is clear that they do not die simultaneously in the CSBP time scale, since their death time in this time scale is simply their height. As U is a continuous mapping, we deduce that they do not die simultaneously in the coalescent time scale either. $\square$

The class of measures Λ concerned by Theorem 1 can be enlarged a little thanks to the following corollary.

Corollary 4.5 *Let Λ be a finite measure such that $x^{-2}\Lambda(dx) = x^{-2}\text{Beta}(2 - \alpha, \alpha)(dx) + \mu(dx)$ where μ is a finite measure on $(0, 1)$ and $1 < \alpha \leq 2$, then $\mathbb{P}(E) = 0$ for the corresponding Λ Fleming-Viot process.*

Proof The proof of Lemma 4.2 has shown that $E \in \mathcal{F}_{0+}$. That is, E depends on the very initial behaviour of ρ . Adding a finite mass μ to the measure $x^{-2}\Lambda(dx)$ does not change the initial behaviour of the corresponding Λ Fleming-Viot process. Indeed, in the lookdown representation, the first birth event induced by μ arrives at a time T , distributed as an exponential random variable with parameter $\mu((0, 1))$, which is strictly positive almost surely. Hence until time T , the behaviour of the Λ Fleming-Viot process coincides with the behaviour of the $\text{Beta}(2 - \alpha, \alpha)$ Fleming-Viot process. Then, the previous proposition ensures the asserted result. $\square$

Finally, we conjecture that this result is true for any Λ coalescent that comes down from infinity.

Conjecture 4.6 *If $\Lambda \in \text{CDI}$, $\mathbb{P}(E) = 0$.*

4.2 An ordering of the ancestral types

The goal of this subsection is to present an ordering of the ancestral types, which is consistent with the definition of the primitive Eve of Bertoin and Le Gall as explained in the introduction. We treat two cases: **[(CDI) and $\mathbb{P}(E) = 0$]** and **[Bolthausen-Sznitman]**.

In the **[(CDI) and $\mathbb{P}(E) = 0$]** case, almost surely the process $t \mapsto \#\rho_t$ is non-increasing, makes negative jumps of size 1, starts from $+\infty$ and reaches 1 in finite time. Denote by d^i the jump time of this process from i to $i - 1$ and let $\mathbf{e}^i$ be the ancestral type that gets extinct at this time, for every $i \geq 2$. Additionally, let $\mathbf{e}^1$ be the primitive Eve defined by Bertoin and Le Gall. Then we obtain a reordering $(\mathbf{e}^i)_{i \geq 1}$ of the ancestral types.

In the **[Bolthausen-Sznitman]** case, we let $\mathbf{e}^1$ be the primitive Eve defined by Bertoin and Le Gall, that is, the random point in $[0, 1]$ such that

$$\rho_t(\{\mathbf{e}^1\}) \xrightarrow[t \uparrow \infty]{} 1$$

This proposition, whose proof is given in Section 8, exhibits a reordering of the ancestral types.

Proposition 4.7 *Almost surely and recursively for each integer $i > 1$, there exists a r.v. $\mathbf{e}^i$ such that*

$$\lim_{t \rightarrow \infty} \frac{\rho_t(\{\mathbf{e}^i\})}{1 - \sum_{1 \leq j \leq i-1} \rho_t(\{\mathbf{e}^j\})} = 1 \quad (32)$$

Remark 4.8 *The proof of this proposition makes use of the connection with continuous-state branching processes in the Bolthausen-Sznitman case [5]. This connection can be used in the α -stable case with $\alpha \in (0, 1)$ to prove that for any two ancestral types x, y distinct from the primitive Eve the ratio $\frac{\rho_t(\{x\})}{\rho_t(\{y\})}$ converges to a value in $(0, \infty)$. Therefore one cannot order the ancestral types in a consistent manner for those measures Λ .*

In both cases, we have defined an ordering $(\mathbf{e}^i)_{i \geq 1}$ of the ancestral types : we call this sequence the *Eves*. We now give several results that prove that this ordering is relevant. The following proposition shows that if there exists a lookdown representation whose process of limiting empirical measures is equal to the process ρ , then the sequence of initial types of the lookdown is necessarily our ordering of the ancestral types of ρ .

Proposition 4.9 *Consider a Λ flow of partitions $\hat{\Pi}$ and a sequence $(\chi_0(i))_{i \geq 1}$ of r.v. taking distinct values in $[0, 1]$. Let $(\Xi_t, t \in [0, \infty)) := \mathcal{E}_0(\hat{\Pi}, (\xi_0(i))_{i \geq 1})$ be the limiting empirical measures of the lookdown process defined from these objects. If $(\Xi_t, t \in [0, \infty)) = (\rho_t, t \in [0, \infty))$ a.s., then $(\xi_0(i))_{i \geq 1} = (\mathbf{e}^i)_{i \geq 1}$ a.s.*

Remark 4.10 *Here the trajectories of the flow of partitions are assumed to be deterministic flows of partitions. See Subsection 3.2.*

Proof Denote by Ω^* the event of probability 1 on which $(\Xi_t, t \in [0, \infty)) = (\rho_t, t \in [0, \infty))$. We know that there exists a random permutation σ of $\mathbb{N}$ such that on Ω^*

$$(\xi_0(i))_{i \geq 1} = (\mathbf{e}_0^{\sigma(i)})_{i \geq 1}$$

To prove the proposition, we have to show that, on Ω^* , σ is the identical permutation of $\mathbb{N}$. We implicitly restrict ourselves to this event for the rest of the proof.

[(CDI) and $\mathbb{P}(E) = 0$] case. It suffices to remark that both $(\mathbf{e}_0^i)_{i \leq 1}$ and $(\xi_0(i))_{i \geq 1}$ are ordered by decreasing persistence. Indeed, in the lookdown representation, the lower an initial type is located (in the set of levels) the longer it will survive. We deduce that σ is the identical permutation.

[Bolthausen-Sznitman] case. The points $(\mathbf{e}^i)_{i \geq 1}$ are ordered by decreasing masses at ∞ , that is, for all $i \geq 1$

$$\frac{\rho_t(\mathbf{e}^i)}{\sum_{j \geq i} \rho_t(\mathbf{e}^j)} \xrightarrow[t \rightarrow \infty]{} 1 \quad (33)$$

Thus it implies that for all $i \geq 1$ we have

$$\mathbb{P}[(\rho_t(\mathbf{e}^j))_{1 \leq j \leq i} = (\rho_t(\mathbf{e}^j))_{1 \leq j \leq i}^\downarrow] \xrightarrow[t \rightarrow \infty]{} 1$$

where symbol $\downarrow$ is the reordering operator of the corresponding sequence by decreasing values. Now remark that for all $t \geq 0$ we have

$$(\rho_t(\mathbf{e}^i))_{i \geq 1}^\downarrow = (\Xi_t(\xi_0(i)))_{i \geq 1}^\downarrow \quad (34)$$

Thanks to the lookdown construction, we know that $\Xi_t(\xi_0(i)) = |\hat{\Pi}_{0,t}(i)|$ for all $t \geq 0$ almost surely. From the exchangeability of $\hat{\Pi}_{0,t}$, we deduce that the sequence $(\Xi_t(\xi_0(i)))_{i \geq 1}$ is a size-biased reordering of $(\Xi_t(\xi_0(i)))_{i \geq 1}^\downarrow$ (see Section 2.1.3 in [4]). But the mass-partition $(\Xi_t(\xi_0(i)))_{i \geq 1}^\downarrow$ has a particular form thanks to Equations (33) and (34). We deduce that for every $i \geq 1$

$$\mathbb{P}[(\Xi_t(\xi_0(j)))_{1 \leq j \leq i} = (\Xi_t(\xi_0(j)))_{1 \leq j \leq i}^\downarrow] \xrightarrow{t \rightarrow \infty} 1$$

So we get for every $i \geq 1$

$$\mathbb{P}[(\Xi_t(\xi_0(i))) = (\rho_t(\mathbf{e}_0^i))] \xrightarrow{t \rightarrow \infty} 1$$

Therefore, this last identity together with the fact that two distinct ancestral types have two distinct asymptotic behaviours imply that $\xi_0(i) = \mathbf{e}^i$ for every $i \geq 1$. $\square$

We now determine the distribution of the sequence $(\mathbf{e}^i)_{i \geq 1}$. In addition, we prove that this sequence of Eves is independent with its process of asymptotic frequencies. This result allows one to decompose a Λ Fleming-Viot process into two independent objects : the Eves and its process of asymptotic frequencies. The dynamic version of this decomposition is provided in the next section.

Proposition 4.11 *The sequence $(\mathbf{e}^i)_{i \geq 1}$ is a sequence of i.i.d. $\text{uniform}[0, 1]$ r.v. independent of the sequence of processes $(\rho_t(\mathbf{e}^i), t \in [0, \infty))_{i \geq 1}$.*

Proof Denote by Φ the measurable map that associates to a Λ Fleming-Viot process its ancestral types according to the definition given at the previous subsection. In particular, we have $\Phi(\rho) = (\mathbf{e}^i)_{i \geq 1}$. Now consider a sequence $(\xi_0(i))_{i \geq 1}$ of i.i.d. $\text{uniform}[0, 1]$ r.v., and an independent Λ flow of partitions $\hat{\Pi}$ (with trajectories assumed to be deterministic flows of partitions). Denote by $(\Xi_t, t \geq 0) = \mathcal{E}_0(\hat{\Pi}, (\xi_0(i))_{i \geq 1})$ the limiting empirical measures of the lookdown process defined from the latter objects. As recalled in Subsection 2.3, $(\Xi_t, t \geq 0)$ is a Λ Fleming-Viot process. Hence we can define its ancestral types $\Phi(\Xi)$. From Proposition 4.9, we deduce that a.s.

$$\Phi(\Xi) = (\xi_0(i))_{i \geq 1}$$

Therefore, using the fact that $(\rho_t, t \geq 0) \stackrel{(d)}{=} (\Xi_t, t \geq 0)$, we deduce that

$$(\rho, \Phi(\rho)) \stackrel{(d)}{=} (\Xi, \Phi(\Xi)) \quad (35)$$

This implies that $(\mathbf{e}_0^i)_{i \geq 1} \stackrel{(d)}{=} (\xi_0(i))_{i \geq 1}$, and thus, it is a sequence of i.i.d. $\text{uniform}[0, 1]$ r.v. Moreover, note that the asymptotic frequencies $(\Xi_t(\xi_0(i)), t \geq 0)_{i \geq 1}$ only depend on the flow of partitions $\hat{\Pi}$, thus are independent of the initial types $(\xi_0(i))_{i \geq 1}$. Using a similar identity in law as in Equation (35), we deduce that the sequence $(\mathbf{e}_0^i)_{i \geq 1}$ is independent of $(\rho_t(\mathbf{e}_0^i), t \in [0, \infty))_{i \geq 1}$. $\square$

5 Eves process and stochastic flow of partitions

In this section, we consider a Λ flow of bridges $(F_{s,t}, -\infty < s \leq t < \infty)$. Recall that for each $s \in \mathbb{R}$, $(\rho_{s,t}, t \in [s, \infty))$ is a càdlàg Λ Fleming-Viot process (we use a modification of the initial flow of bridges, as explained in Subsection 2.2). As in the previous subsection, we treat two cases: **[(CDI) and $\mathbb{P}(E) = 0]$** and **[Bolthausen-Sznitman]**. The dynamic version of the sequence of Eves will be called the Eves process. We will provide a striking result on the coupling of two-dimensional marginals of this process that will allow us to introduce a collection of random partitions $(\hat{\Pi}_{s,t}, s \leq t)$. Finally, we will prove that the latter is a stochastic flow of partitions (Theorem 2).

5.1 The Eves process

First, let us recall the dynamics of the primitive Eve introduced by Bertoin and Le Gall in Section 5.3 of [6]. For every $s \in \mathbb{R}$, let $\mathbf{e}_s^1$ be the r.v. in $[0, 1]$ such that

$$\rho_{s,t}(\{\mathbf{e}_s^1\}) \xrightarrow[t \uparrow \infty]{} 1$$

For each $s \in \mathbb{R}$, the definition holds on an event of probability 1. On the complementary event, set $\mathbf{e}_s^1 := 0$.

From the very definition of $\mathbf{e}_s^1$, one has $\mathbf{e}_s^1 \in [F_{0,s}(\mathbf{e}_0^1-), F_{0,s}(\mathbf{e}_0^1)]$ thanks to the cocycle property. This implies the following result due to Bertoin and Le Gall [6], Section 5.3.

Proposition 5.1 *For all $s > 0$, almost surely $\mathbf{e}_0^1 = F_{0,s}^{-1}(\mathbf{e}_s^1)$.*

This identity describes the backward evolution of the primitive Eve process. We now define the Eves process and will see in the next subsection that a similar identity holds for this process.

Fix a time $s \in \mathbb{R}$, and consider the Λ Fleming-Viot $(\rho_{s,t}, t \in [s, \infty))$. We define the sequence $(\mathbf{e}_s^i)_{i \geq 1}$ using the definition of Eves of Subsection 4.2 applied to $(\rho_{s,t}, t \in [s, \infty))$. Remark that this definition holds on an event Ω_s of probability 1. On the event $\Omega \setminus \Omega_s$ of zero probability, we set any arbitrary values to the sequence $(\mathbf{e}_s^i)_{i \geq 1}$.

Notice from the definition, that for any fixed time s , the sequence $(\mathbf{e}_s^i)_{i \geq 1}$ is independent of the past of the flow of bridges until time s . Indeed, this sequence only depends on bridges of the form $F_{s,t}, t \geq s$, and thanks to the independence property of the increments of a flow of bridges, we obtain the claimed independence. In the next subsection, we provide a generalization of Proposition 5.1.

5.2 Key property and the flow of partitions

Consider the Eves at two distinct times : $(\mathbf{e}_s^i)_{i \geq 1}$ and $(\mathbf{e}_0^j)_{j \geq 1}$. Our next proposition provides the relation between those two sequences, and refers to a key property used by Bertoin and Le Gall in Lemma 2 in [6] which we now recall. Consider a sequence of independent uniform $[0, 1]$ variables $(V_i)_{i \geq 1}$ and an independent bridge B . Denote by $(A_j)_{j \geq 1}$ the blocks of the partition $\pi(B, (V_i)_{i \geq 1})$ ordered by their smallest element (those blocks are in finite number if B has a finite number of jumps and no drift). Define

a sequence of random variables $V'_j := B^{-1}(V_i)$ for an arbitrary $i \in A_j$. If there is a finite number of blocks in $\pi(B, (V_i)_{i \geq 1})$, complete the sequence with independent uniform $[0, 1]$ random variables. The key property yields that the $(V'_j)_{j \geq 1}$ are i.i.d uniform $[0, 1]$ variables, independent of $\pi(B, (V_i)_{i \geq 1})$. We will say that $(B, (V_i)_{i \geq 1}, (V'_j)_{j \geq 1})$ follows the *composition rule*.




Figure 2: An illustration of Proposition 5.2 in the (CDI) case. On the left, an example of the composition rule. On the right, the genealogical structure arising from this result. The corresponding partition (restricted to $[4]$) is $\hat{\Pi}_{0,s} = \{\{1, 3\}, \{2\}, \{4\}\}$.

Proposition 5.2 *For all $s > 0$, almost surely $(F_{0,s}, (e_s^i)_{i \geq 1}, (e_0^j)_{j \geq 1})$ follows the composition rule.*

The following technical proof is illustrated by Figure 2.

Proof There exists an event $\tilde{\Omega} \subset \Omega_0 \cap \Omega_s$ of probability 1 such that on this event for all $t \in \mathbb{Q}$, $F_{0,t} = F_{s,t} \circ F_{0,s}$. We work on this event until the end of the proof. Recall that $(e_s^i)_{i \geq 1}$ is a sequence of i.i.d uniform $[0, 1]$ r.v. independent of the bridge $F_{0,s}$. Those r.v. play the role of the $(V_i)_{i \geq 1}$ in the key property presented above.

Let K be the random number of blocks of $\pi := \pi(F_{0,s}, (e_s^i)_{i \geq 1})$. This number is finite in the **[(CDI) and $\mathbb{P}(E) = 0$]** case while it is infinite in the **[Bolthausen-Sznitman]** case, almost surely. Denote by $(A_j)_{1 \leq j \leq K}$ the blocks in the increasing order of their least element. Then, we can define a sequence of random variables $V'_j := F_{0,s}^{-1}(e_s^{i_j})$ where $i_j := \min(A_j)$, for all $j \in [K]$. When K is finite, we set $V'_j := e_0^j$ for all $j > K$. The key property of Bertoin and Le Gall ensures that the $(V'_j)_{j \in [K]}$ are independent of the partition π . To prove the proposition, it remains to show that:

- (i) $(e_0^j)_{j > K}$ are i.i.d. uniform $[0, 1]$, independent of $(e_0^j)_{j \in [K]}$ and of π .
- (ii) $e_0^j = V'_j$ for all $j \in [K]$ a.s.

We start with the first assertion. In the **[Bolthausen-Sznitman]** case, this assertion is trivial since $K = \infty$ a.s. We consider the **[(CDI) and $\mathbb{P}(E) = 0$]** case. Since K only depends on the values of the sequence $(\rho_{0,s}(\{e_0^i\}))_{i \geq 1}$, we deduce from Proposition 4.11 that $(e_0^j)_{j > K}$ are i.i.d. uniform $[0, 1]$, independent of

$(\mathbf{e}_0^j)_{j \in [K]}$. Furthermore, since $F_{0,s}$ only depends on $(\mathbf{e}_0^j)_{j \in [K]}$ and on $(\rho_{0,s}(\{\mathbf{e}_0^i\}))_{i \in [K]}$, we deduce that the $(\mathbf{e}_0^j)_{j > K}$ are independent of $F_{0,s}$. Finally since $(\mathbf{e}_0^j)_{j > K}$ are independent of the future of the flow F after time s , it is clear that they are independent of $(\mathbf{e}_s^i)_{i \geq 1}$. Therefore, they are independent of both $(\mathbf{e}_s^i)_{i \geq 1}$ and $F_{0,s}$, and so, of π . The first assertion follows. Let us prove the second assertion.

Note that the $(\mathbf{e}_0^j)_{j \in [K]}$ are a reordering of the $(V_j')_{j \in [K]}$. Indeed, the $(\mathbf{e}_0^j)_{j \in [K]}$ correspond to the jump locations of $F_{0,s}$, and the $(V_j')_{j \in [K]}$ form the set of values taken by $(F_{0,s}^{-1}(\mathbf{e}_s^i))_{i \geq 1}$.

We stress that for each $j \in [K]$, for all $t \in [s, \infty)$ (see Figure 2)

$$\rho_{s,t}(\{\mathbf{e}_s^{i_j}\}) \leq \rho_{0,t}(\{V_j'\}) \leq \rho_{s,t}([0,1] \setminus \{\mathbf{e}_s^1, \dots, \mathbf{e}_s^{i_j-1}\})$$

Let us give an informal argument for this inequality. V_j' is a type at time 0 that carries a positive fraction of the population at time s , this fraction of the population includes the individual $\mathbf{e}_s^{i_j}$ but none of the individuals $\{\mathbf{e}_s^1, \dots, \mathbf{e}_s^{i_j-1}\}$. When considering the descences of these individuals at time $t \geq s$, we see that the descence of $\mathbf{e}_s^{i_j}$ is included into the descence of V_j' , and therefore in the descence of $[0,1] \setminus \{\mathbf{e}_s^1, \dots, \mathbf{e}_s^{i_j-1}\}$. The inequality follows by taking asymptotic frequencies.

Case 1 : CDI and $\mathbb{P}(E) = 0$ Both processes $t \mapsto \rho_{s,t}([0,1] \setminus \{\mathbf{e}_s^1, \dots, \mathbf{e}_s^{i_j-1}\})$ and $t \mapsto \rho_{s,t}(\{\mathbf{e}_s^{i_j}\})$ reach 0 at the same time $d_s^{i_j} := \inf\{t \geq s : \rho_{s,t}(\{\mathbf{e}_s^{i_j}\}) = 0\}$, so we conclude that the same holds for the process $t \mapsto \rho_{0,t}(\{V_j'\})$. Hence, the collection of points $(V_j')_{j \in [K]}$ is ordered by decreasing persistence. Since the $(\mathbf{e}_0^j)_{j \geq 1}$ are a reordering of the $(V_j')_{j \geq 1}$, we conclude that $V_j' = \mathbf{e}_0^j$ for all $j \in [K]$.

Case 2 : Bolthausen-Sznitman We know that :

$$\lim_{t \rightarrow \infty} \rho_{0,t}(\{F_{0,s}^{-1}(\mathbf{e}_s^1)\}) = 1 \quad (36)$$

By definition of the Eves process, it follows that $\mathbf{e}_0^1 = V_1'$.

From Proposition 4.7 we know that for each $j \geq 2$

$$\lim_{t \rightarrow \infty} \frac{\rho_{s,t}(\{\mathbf{e}_s^{i_j}\})}{1 - \sum_{1 \leq l \leq i_j-1} \rho_{s,t}(\{\mathbf{e}_s^l\})} = 1 \quad (37)$$

thus we get

$$\lim_{t \rightarrow \infty} \frac{\rho_{0,t}(\{V_j'\})}{1 - \sum_{1 \leq l \leq j-1} \rho_{0,t}(\{V_l'\})} \geq \lim_{t \rightarrow \infty} \frac{\rho_{s,t}(\{\mathbf{e}_s^{i_j}\})}{1 - \sum_{1 \leq l \leq i_j-1} \rho_{s,t}(\{\mathbf{e}_s^l\})} = 1 \quad (38)$$

Since the $(\mathbf{e}_0^j)_{j \geq 1}$ are a reordering of the $(V_j')_{j \geq 1}$, we deduce from the previous inequation and Proposition 4.7 that $V_j' = \mathbf{e}_0^j$ for each $j \geq 2$.

This ends the proof of the proposition. $\square$

Remark 5.3 In the [(CDI) and $\mathbb{P}(E) = 0$] case, the number of blocks K of the partition $\pi(F_{0,s}, (\mathbf{e}_s^i)_{i \geq 1})$ is finite a.s. The descendents of individuals $(\mathbf{e}_0^j)_{j > K}$ have become extinct by time s . In the genealogical interpretation, $(\mathbf{e}_0^j)_{1 \leq j \leq K}$ are the ancestors of the K oldest families in the population alive at time s .

In the setting of flow of bridges, the population model has been complexified: in particular, $\mathbf{e}_s^i$ is jointly seen as a type and an individual. Let us clarify this interpretation. At each time $s \in \mathbb{R}$, we consider an infinite population where each individual is characterized by a type in $[0, 1]$. This population gives birth to a descendance at time $s+$, and therefore genealogical relationships exist between this population at time s and those at time $t > s$. The Eves $(\mathbf{e}_s^i)_{i \geq 1}$ identify those individuals who take a significant part to this genealogical structure and are seen as the ancestors at time s of a positive fraction of the population at a time $t > s$. To characterize this ancestor, we use a type between $[0, 1]$ given by the point $\mathbf{e}_s^i$. Among its descendents at time $t > s$, we identify those who will have a significant role in the genealogical structure after time t . Note that the descendents of $\mathbf{e}_s^i$ do not have necessarily the same type as their ancestor.

The preceding proposition motivates the introduction of the following random partition

$$\hat{\Pi}_{s,t} := \pi(F_{s,t}, (\mathbf{e}_t^i)_{i \geq 1}) \quad (39)$$

for all $s \leq t$. Indeed this proposition has shown that the i -th block of $\hat{\Pi}_{s,t}$ corresponds to the descendents of $\mathbf{e}_s^i$ in the population at time t , for each $i \geq 1$ (see Figure 2 for an illustration). Thus this partition provides the genealogical relationships between the Eves.

Theorem 2 *The collection of partitions $(\hat{\Pi}_{s,t}, -\infty < s \leq t < \infty)$ is a Λ flow of partitions.*

This theorem somehow extends the basic correspondence of Bertoin and Le Gall, Section 3.2 in [6]. Moreover, this is a pathwise correspondence since the flow of partitions has been defined in terms of the flow of bridges.

Proof The first requirement of Definition 3.6 is a consequence of Proposition 5.2. Indeed, the partition $\hat{\Pi}_{r,t}$ is obtained by applying the key property with the sequence $(\mathbf{e}_t^i)_{i \geq 1}$ and the bridge $F_{r,t} = F_{s,t} \circ F_{r,s}$ a.s. From the proof of Corollary 1, in [6], we deduce that the random partition $\hat{\Pi}_{r,t}$ is equal to the coagulation of $\hat{\Pi}_{s,t}$ by the partition $\hat{\Pi}_{r,s}$ (although this corollary asserts an equality in distribution, the proof actually defines an a.s. equality).

Let us prove the independence of the increments in the case $n = 2$, the general case is obtained by an easy induction. Fix $r < s < t$. Thanks to Proposition 5.2, we know that the sequence $(\mathbf{e}_s^i)_{i \geq 1}$ is independent of the partition $\hat{\Pi}_{s,t}$. Additionally, the bridge $F_{r,s}$ is independent of the partition $\hat{\Pi}_{s,t}$ from the independence of the increments of a flow of bridges and the definition of the latter partition. Given that the partition $\hat{\Pi}_{r,s}$ depends only on the sequence $(\mathbf{e}_s^i)_{i \geq 1}$ and the bridge $F_{r,s}$, we deduce the independence of $\hat{\Pi}_{r,s}$ and $\hat{\Pi}_{s,t}$. Furthermore, the fact that the distribution of $\hat{\Pi}_{s,t}$ only depends on $t - s$ is an immediate consequence of the stationarity of flows of bridges.

The convergence in probability of $\hat{\Pi}_{0,t} \rightarrow 0_{[\infty]}$ for the distance $d_{\mathcal{P}}$ is a consequence of the next lemma. Finally, since the flow of bridges F is associated with the measure Λ , we immediately deduce that $(\hat{\Pi}_{-t,0}, t \geq 0)$ is a Λ coalescent using the result of Bertoin and Le Gall recalled in Subsection 2.2. $\square$

Lemma 5.4 Consider a collection of bridges $(B_t)_{t>0}$ and an independent sequence of i.i.d. uniform $[0, 1]$ random variables $(V_i)_{i\geq 1}$. The following conditions are equivalent

- a) The exchangeable partition $\pi(B_t, (V_i)_{i\geq 1})$ converges in probability to $0_{[\infty]}$ for the distance $d_{\mathcal{P}}$ as $t \downarrow 0$.
- b) The bridge B_t converges in probability to Id in the sense of Skorohod's topology as $t \downarrow 0$.

We postpone the proof of this lemma to Section 8.

6 Proof of Theorem 3

We have defined a stochastic flow of partitions $\hat{\Pi}$ pathwise from the flow of bridges F . As F may have many irregularities, so may $\hat{\Pi}$. However, using the regularization procedure described in Subsection 3.2, one obtains a modification of the original flow $\hat{\Pi}$, still denoted $\hat{\Pi}$ for simplicity, such that its trajectories are deterministic flows of partitions, almost surely. Then, the collection of the jumps of $\hat{\Pi}$ defines a Λ lookdown graph as proved in Proposition 3.14. Thus, we have defined pathwise from the flow of bridges F an Eves process $(\mathbf{e}_s^i, s \in \mathbb{R})_{i\geq 1}$ and a lookdown graph. The goal of this section is to prove that those two objects define a collection of coupled lookdown processes with limiting empirical measures the collection $(\rho_{s,t}, t \in [s, \infty))_{s \in \mathbb{R}}$.

Let us introduce a particle system $(\xi_{s,t}(i), s \leq t)_{i\geq 1}$ as follows. For each $s \in \mathbb{R}$, set $(\xi_{s,t}(i), t \in [s, \infty))_{i\geq 1} := \mathcal{L}_s(\hat{\Pi}, (\mathbf{e}_s^i)_{i\geq 1})$ and denote by $(\Xi_{s,t}, t \in [s, \infty)) := \mathcal{E}_s(\hat{\Pi}, (\mathbf{e}_s^i)_{i\geq 1})$ its process of limiting empirical measures. Let us recall the statement of the theorem.

Theorem 3 The flow of Λ Fleming-Viot processes is uniquely decomposed into two random objects : the flow of partitions and the Eves process. More precisely, for each $s \in \mathbb{R}$ almost surely

$$\mathcal{E}_s(\hat{\Pi}, (\mathbf{e}_s^i)_{i\geq 1}) = (\rho_{s,t}, t \in [s, \infty))$$

Furthermore if $\hat{\Pi}'$ is a Λ flow of partitions and for each $s \in \mathbb{R}$, $(\chi_s(i))_{i\geq 1}$ is a sequence of r.v. taking distinct values in $[0, 1]$ such that we have almost surely $\mathcal{E}_s(\hat{\Pi}', (\chi_s(i))_{i\geq 1}) = (\rho_{s,t}, t \in [s, \infty))$ then

- For each $s \in \mathbb{R}$, a.s. $(\chi_s(i))_{i\geq 1} = (\mathbf{e}_s^i)_{i\geq 1}$.
- Almost surely, $\hat{\Pi}' = \hat{\Pi}$.

This section is devoted to the proof of the theorem. In the first subsection, we prove the decomposition statement. In the second subsection, we investigate the uniqueness properties of the lookdown construction and prove the uniqueness statement. Finally, in the third subsection we compare our lookdown construction from a flow of bridges with the original lookdown definition of Donnelly and Kurtz in [12].

6.1 Decomposition

Proof (Theorem 3-Decomposition) Fix $s \in \mathbb{R}$. Remark that both processes $(\Xi_{s,t}, t \in [s, \infty))$ and $(\rho_{s,t}, t \in [s, \infty))$ are càdlàg processes. Therefore, to prove that a.s.

$$(\Xi_{s,t}, t \in [s, \infty)) = (\rho_{s,t}, t \in [s, \infty))$$

it is sufficient to prove that for each $t \in [s, \infty)$, we have a.s.

$$\Xi_{s,t} = \rho_{s,t}$$

Consider a time $t \in [s, \infty)$. Since $\hat{\Pi}_{s,t}$ is exchangeable it admits asymptotic frequencies. A simple application of Equations (14) and (20) ensures that

$$\Xi_{s,t}(dx) = \sum_{i=1}^{\infty} |\hat{\Pi}_{s,t}(i)| \delta_{\mathbf{e}_s^i}(dx)$$

Moreover, thanks to Proposition 5.2 we know that a.s. for every $i \geq 1$, $\rho_{s,t}(\{\mathbf{e}_s^i\}) = |\hat{\Pi}_{s,t}(i)|$. Therefore, a.s.

$$\rho_{s,t}(dx) = \sum_{i=1}^{\infty} |\hat{\Pi}_{s,t}(i)| \delta_{\mathbf{e}_s^i}(dx)$$

Thus, we get that a.s. for every $t \in [s, \infty) \cap \mathbb{Q}$, we have $\Xi_{s,t}(\cdot) = \rho_{s,t}(\cdot)$. Since both are càdlàg processes, we have proved the identity. This ensures the decomposition statement of the theorem. $\square$

6.2 Uniqueness

We now focus on the uniqueness statement of the theorem. Let $\hat{\Pi}'$ be a Λ flow of partitions. We assume that the trajectories of $\hat{\Pi}'$ are deterministic flows of partitions, otherwise the lookdown construction could not be applied to this object. But from the regularization results of Subsection 3.2, this assumption can be made without loss of generality. For each $s \in \mathbb{R}$, consider a sequence $(\chi_s(i))_{i \geq 1}$ of r.v. taking distinct values in $[0, 1]$. We denote by $(X_{s,t}, t \in [s, \infty)) := \mathcal{E}_s(\hat{\Pi}', (\chi_s(i))_{i \geq 1})$ the process of limiting empirical measures of the corresponding lookdown process. We suppose that for each $s \in \mathbb{R}$, almost surely $(X_{s,t}, t \in [s, \infty)) = (\rho_{s,t}, t \in [s, \infty))$.

Proof (Theorem 3-Uniqueness) Proposition 4.9 implies that for each $s \in \mathbb{R}$, we have $(\chi_s(i))_{i \geq 1} = (\mathbf{e}_s^i)_{i \geq 1}$ a.s. It remains to prove that $\hat{\Pi}' = \hat{\Pi}$ a.s.

There exists an event Ω^* of probability 1 such that on this event, for every rational values $s \leq t$ and every integer $i \geq 1$ we have

$$\rho_{s,t}(\{\mathbf{e}_s^i\}) = |\hat{\Pi}_{s,t}(i)| = |\hat{\Pi}'_{s,t}(i)| \quad (40)$$

Some properties will hold both for $\hat{\Pi}'$ and $\hat{\Pi}$, thus we will use the notation $\hat{\Pi}^\times$ to designate indifferently any of them. The proof of the uniqueness statement of the theorem relies on the following result.

Proposition 6.1 *The knowledge of $(|\hat{\Pi}_{s,t}^\times(i)|, s \leq t \in \mathbb{Q})_{i \geq 1}$ is sufficient to recover the flow of partitions $(\hat{\Pi}_{s,t}^\times, -\infty < s \leq t < \infty)$.*

Applying this proposition to both flows $\hat{\Pi}$ and $\hat{\Pi}'$ together with Equation (40), we obtain that $\hat{\Pi} = \hat{\Pi}'$ almost surely. $\square$

It remains to prove Proposition 6.1. To that end, we remark that a flow of partitions is completely characterized by its jumps $\hat{\Pi}_{s-,s}^\times$ for all $s \in \mathbb{R}$. From Proposition 2.1 (notice that in this proposition, the coalescent is càdlàg while in the flow of partitions the coalescent is làdcàg but this does not induce any difficulty), we deduce that almost surely for all $s \in \mathbb{R}$ and all $t \in (s, \infty) \cap \mathbb{Q}$ the quantities $(|\hat{\Pi}_{s-,t}^\times(i)|)_{i \geq 1}$ and $(|\hat{\Pi}_{s,t}^\times(i)|)_{i \geq 1}$ are derived from $(|\hat{\Pi}_{p,q}^\times(i)|, p \leq q \in \mathbb{Q}^2)_{i \geq 1}$ by taking right and left limits. The following lemma asserts that the preceding asymptotic frequencies are sufficient to determine all the jumps $\hat{\Pi}_{s-,s}^\times$, and Proposition 6.1 is proved.

Lemma 6.2 *Let I be a subset of $\mathbb{N}$. The following assertions are equivalent*

i) $\hat{\Pi}_{s-,s}^\times$ has a unique non-singleton block I .

$$ii) \begin{cases} (|\hat{\Pi}_{s-,t}^\times(i)|, t \in (s, \infty) \cap \mathbb{Q}) = (\sum_{j \in I} |\hat{\Pi}_{s,t}^\times(j)|, t \in (s, \infty) \cap \mathbb{Q}) \text{ if } i = \min(I) \\ (|\hat{\Pi}_{s-,t}^\times(i)|, t \in (s, \infty) \cap \mathbb{Q}) = (|\hat{\Pi}_{s,t}^\times(j)|, t \in (s, \infty) \cap \mathbb{Q}) \text{ if } \begin{cases} i \neq \min(I) \\ i = j - (\#\{I \cap [j]\} - 1) \vee 0 \end{cases} \end{cases}$$

In words, this lemma says that we can deduce the jumps $\hat{\Pi}_{s-,s}^\times$ from the evolution of the asymptotic frequencies.

Proof Suppose i). Since $\hat{\Pi}_{s-,t}^\times = Coag(\hat{\Pi}_{s,t}^\times, \hat{\Pi}_{s-,s}^\times)$, the very definition of the coagulation operator implies ii).

Suppose ii). Since the trajectories of $\hat{\Pi}^\times$ are deterministic flows of partitions without simultaneous mergers, we know that $\hat{\Pi}_{s-,s}^\times$ is a partition with at most one non-singleton block.

We know that for all $\omega \in \Omega^*$, $(|\hat{\Pi}_{s-,t}^\times(i)|(\omega), t \in (s, \infty) \cap \mathbb{Q})_{i \geq 1}$ are all distinct. Indeed, in the **[(CDI) and $\mathbb{P}(E) = 0]$** case, the extinction times of the asymptotic frequencies are strictly distinct while in the **[Bolthausen-Sznitman]** case, their asymptotic behaviours are strictly distinct. The same holds for $(|\hat{\Pi}_{s,t}^\times(j)|(\omega), t \in (s, \infty) \cap \mathbb{Q})_{j \geq 1}$. Since $\hat{\Pi}_{s-,t}^\times = Coag(\hat{\Pi}_{s,t}^\times, \hat{\Pi}_{s-,s}^\times)$, the equations of ii) imply that the partition $\hat{\Pi}_{s-,s}^\times$ has a unique non-singleton block I . $\square$

6.3 A remark on the lockdown ordering

Our pathwise lockdown construction from a stochastic flow of bridges is an infinite dimensional extension of the lockdown construction from the Moran model (see [12], where Model I is a Moran model whereas Model II is a lockdown process). Fix $n \in \mathbb{N}$, and consider a Moran model $(x_t(1), \dots, x_t(n), t \geq 0)$ started from a sequence of n i.i.d. uniform $[0, 1]$. Donnelly and Kurtz introduce a lockdown process $(\xi_t(1), \dots, \xi_t(n), t \geq 0)$ on $[0, \infty)$ by defining a random permutation θ of $[n]$ and setting

$$(\xi_t(1), \dots, \xi_t(n)) = (x_t(\theta^{-1}(1)), \dots, x_t(\theta^{-1}(n)))$$

A careful reading of the proof of Theorem 1.1 in [12] shows that θ^{-1} is the random permutation of $[n]$ such that $(x_0(\theta^{-1}(i)))_{i \in [n]}$ is ordered by persistence, that is $x_0(\theta^{-1}(1))$ is the type that fixes, then

$x_0(\theta^{-1}(2))$ is the last type that becomes extinct and so on.

Since a stochastic flow of bridges is somehow an infinite dimensional extension of the Moran model, one should compare Theorem 1.1 of [12] and our lockdown construction (in the **[(CDI) and $\mathbb{P}(E) = 0$]** case). Indeed, both rely on the reordering of the ancestral types by decreasing persistence. Moreover, in both cases, this random reordering is independent of the past of the underlying process (Moran model or flow of bridges). Hence, the random reordering depends on the future of the underlying process and as time passes, the evolution of this process allows one to determine the reordering from the highest levels to the lowest. Note that this implies that the filtration of the Moran model / flow of bridges is not the same as the filtration of the lockdown process.

7 Encoding of the Beta Fleming-Viot process

In this section, we reformulate the results of Berestycki et al. in [1, 2] on the encoding of the lockdown process associated with the $\text{Beta}(2 - \alpha, \alpha)$ Fleming-Viot into a height process associated with an α -stable branching mechanism (α -stable height process for short), with $\alpha \in (1, 2]$, in terms of the flow of partitions and the Eves process. We fix $\alpha \in (1, 2]$ and consider the α -stable branching mechanism $\Psi(q) = q^\alpha$ when $\alpha \in (1, 2)$ and $\Psi(q) = 2q^2$ when $\alpha = 2$. Recall that the notation $\text{Beta}(2 - \alpha, \alpha)$ refers to the measure given by Equation (8) when $\alpha \in (1, 2)$ while it denotes the measure $\delta_0(dx)$ when $\alpha = 2$. Denote by H an α -stable height process as defined in [13] and let ν^{exc} be its excursion measure on $\mathcal{E}_+$, defined to be the set of positive continuous excursions away from 0. Proposition 1.3.3 in [13] ensures the existence of a jointly measurable modification $(L(t, x), t \geq 0, x \geq 0)$ of the local-time accumulated by H at level $x \geq 0$ until time $t \geq 0$, which is continuous in t (the normalization of this local-time is given in that proposition).

Set $T_x := \inf\{t \geq 0 : L(t, x) > 1\}$ for all $x \geq 0$: this is the first time at which the local-time of H at level x is greater than 1. It is well known [5, 9, 13] that

$$x \mapsto Z_x := L(T_0, x) \quad (41)$$

is a continuous-state branching process with branching mechanism Ψ , started from 1. In the sequel, we will consider the process H stopped at T_0 , thus for all $s \geq T_0$, $H_s = 0$.

Let us introduce for each $t \geq 0$, a time-change due to [9]

$$U(t) := \begin{cases} \inf\{s > 0 : \int_0^s \frac{\alpha(\alpha-1)\Gamma(\alpha)}{Z_x^{\alpha-1}} dx > t\} & \text{if } \alpha \in (1, 2) \\ \inf\{s > 0 : \int_0^s \frac{4}{Z_x} dx > t\} & \text{if } \alpha = 2 \end{cases} \quad (42)$$

Then U is an increasing bijective map from $[0, \infty)$ to $[0, S)$, where $S := \sup_{s \geq 0} (H_s)$. This time-change allows to transform a $\text{Beta}(2 - \alpha, \alpha)$ Fleming-Viot process into a continuous-state branching process (see Theorem 1.1 in [9]) where $[0, \infty)$ corresponds to coalescent/Fleming-Viot time-scale while $[0, S)$ stands for the branching process time-scale. In [2], Berestycki et al. showed that the lockdown representation of a $\text{Beta}(2 - \alpha, \alpha)$ Fleming-Viot process could be identified into the excursions of the height

process. We reformulate in terms of flow of partitions this encoding, and we show that the Eves process can be also defined in that setting.

For each $x \in [0, S)$ conditional on Z_x , the excursions of H above level x are distributed according to a Poisson point process on $[0, Z_x] \times \mathcal{E}_+$ with intensity measure $dl \otimes \nu^{exc}$. We denote by $(z_x^i, \epsilon_x^i)_{i \geq 1}$ the set of points of this point process ordered by decreasing height of the excursions, that is, ϵ_x^1 is the highest excursion, ϵ_x^2 is the second highest and so on, while z_x^i is the local-time accumulated by H at level x until the beginning of the excursion ϵ_x^i , for each $i \geq 1$.

The Eves are actually related to the local-time locations as follows. For all $s \in [0, \infty)$ and $i \in \mathbb{N}$

$$\mathbf{e}_s^i := \frac{z_{U(s)}^i}{Z_{U(s)}} \quad (43)$$

To state a result on this process, we need to introduce a filtration. We denote by $\mathcal{F}_x$ the σ -field that knows everything about the trajectory of H under the value x . A precise definition of such a filtration can be found in [2, 13].

Proposition 7.1 *For each $s \in [0, \infty)$, $(\mathbf{e}_s^i)_{i \geq 1}$ is a sequence of i.i.d. uniform $[0, 1]$, independent of $\mathcal{F}_{U(s)}$.*

Proof We provide a sketch of the proof. Fix $s \in [0, \infty)$. From Itô's excursion theory, we know that $(z_{U(s)}^i, \epsilon_{U(s)}^i)_{i \geq 1}$ are distributed according to a Poisson point process on $[0, Z_{U(s)}] \times \mathcal{E}_+$ with intensity measure $dl \otimes \nu^{exc}$ ordered by decreasing height of the excursions. We deduce that $(z_{U(s)}^i)_{i \geq 1}$ are i.i.d. uniform $[0, Z_{U(s)}]$. Renormalizing by $Z_{U(s)}$, we obtain that $(\mathbf{e}_s^i)_{i \geq 1}$ is i.i.d. uniform $[0, 1]$ independently of $Z_{U(s)}$. The independence property follows from Lemma 17 in [2]. $\square$

The idea is now to remark that the excursions from level y are embedded into those from level $x < y$. More precisely, fix $x < y \in [0, S)$, there exists a unique $k \in \mathbb{N}$ such that the excursion ϵ_y^i is embedded into the excursion ϵ_x^k for a given integer $i \geq 1$. Thus, we define the random partition $\hat{\Pi}_{s,t}$, where $s := U^{-1}(x)$ and $t := U^{-1}(y)$, as follows

$$i \stackrel{\hat{\Pi}_{s,t}}{\sim} j \Leftrightarrow \epsilon_y^i \text{ and } \epsilon_y^j \text{ belong to the same excursion } \epsilon_x^k, \text{ with } k \in \mathbb{N} \quad (44)$$

Proposition 7.2 *The process $(\hat{\Pi}_{s,t}, 0 \leq s \leq t < \infty)$ is a stochastic flow of partitions associated with the measure $\text{Beta}(2 - \alpha, \alpha)(dx)$.*

Proof For all $0 \leq r \leq s \leq t$, the identity $\hat{\Pi}_{r,t} = \text{Coag}(\hat{\Pi}_{s,t}, \hat{\Pi}_{r,s})$ is an immediate consequence of the definition of the partitions. Moreover we deduce from Proposition 2.1 in [1] and Theorem 1 in [2] that $(\hat{\Pi}_{t-t',t}, t' \in [0, t])$ is a $\text{Beta}(2 - \alpha, \alpha)$ coalescent restricted to $[0, t]$ (recall that $\alpha = 2$ corresponds to the Kingman coalescent). Thus, the law of $\hat{\Pi}_{s,t}$ only depends on $t - s$ and $\hat{\Pi}_{t-r,t} \rightarrow 0_{[\infty]}$ as $r \downarrow 0$ in probability. Furthermore, we stress that for any $0 \leq s < t \leq t_1 < \dots < t_n$, $\hat{\Pi}_{s,t}$ is independent of the partitions $(\hat{\Pi}_{t_i, t_{i+1}})_{i \in [n-1]}$. Indeed, $\hat{\Pi}_{s,t}$ only depends on $\mathcal{F}_{U(t)}$ and on the $(z_{U(t)}^i)_{i \geq 1}$. Thus, conditional

on $Z_{U(t)}$, it is independent of the $(\epsilon_{U(t)}^i)_{i \geq 1}$. Now remark that the partitions $(\hat{\Pi}_{t_i, t_{i+1}})_{i \in [n-1]}$ are defined exclusively from the knowledge of $(\epsilon_{U(t)}^i)_{i \geq 1}$ and that their distribution does not depend on $Z_{U(t)}$. Thus, we deduce that these partitions are independent of $\hat{\Pi}_{s,t}$. $\square$

We have all the elements to define a lookdown process from any given time $s \in [0, \infty)$. Let $(\xi_{s,t}(i), t \in [s, \infty))_{i \geq 1} := \mathcal{L}_s(\hat{\Pi}, (\mathbf{e}_s^i)_{i \geq 1})$ and $(\Xi_{s,t}, t \in [s, \infty)) := \mathcal{E}_s(\hat{\Pi}, (\mathbf{e}_s^i)_{i \geq 1})$. Here, we use directly $\hat{\Pi}$ as a lookdown graph assuming that it has been regularized.

Proposition 7.3 *For all $s \in [0, \infty)$, $(\Xi_{s,t}, t \in [s, \infty))$ is a $\text{Beta}(2 - \alpha, \alpha)$ Fleming-Viot process.*

Proof This follows from the distribution of the Eves and the flow of partitions, together with the lookdown representation. $\square$

One should remark that the mass $\Xi_{s,t}(\mathbf{e}_s^i) = |\hat{\Pi}_{s,t}(i)|$ is exactly the ratio of the local-time accumulated at level $U(t)$ by the excursion $\epsilon_{U(s)}^i$ over the local-time accumulated at this same level by the entire process H , for each $i \geq 1$.

8 Appendix

8.1 Proof of Proposition 2.1

We know that for every $t \geq 0$, almost surely the blocks of Π_t admit asymptotic frequencies by exchangeability. Therefore, this holds simultaneously for all $t \in \mathbb{Q}_+$ on an event Ω^* of probability 1. We now prove that it actually holds simultaneously for all $t \geq 0$ on Ω^* .

We distinguish three cases. First consider the case (CDI). The blocks are finitely many at any positive time and they evolve at discrete times. Then, for any irrational value $t > 0$, there exists two rational values $p < t < q$ such that $\Pi_p = \Pi_{t-}$ and $\Pi_t = \Pi_q$. The result follows.

Second consider the case where the coalescent has dust. Pitman in [17] proved this is equivalent with $\int_0^1 x^{-1} \Lambda(dx) < \infty$. Therefore, for each $i \geq 1$, the number of coalescence events in which the i -th block is involved is finite on any compact interval of time. Thus, for any irrational value $t > 0$, there exists two rational values $p < t < q$ such that $\Pi_p = \Pi_{t-}$ and $\Pi_t = \Pi_q$. This case is complete.

Finally, consider the remaining case : infinitely many blocks without dust. Let t be an irrational positive value and p be any rational positive value lower than t . We have for every $i \geq 1$ (with an obvious notation)

$$\Pi_t(i) = \bigcup_{j \in \Pi_{p,t}(i)} \Pi_p(j)$$

Fix $\epsilon > 0$. As there is no singletons, there exists $k(\omega) \geq 1$ such that $\sum_{j=1}^k \Pi_p(j) > 1 - \epsilon$ and thus

$$\lim_{n \rightarrow \infty} \frac{1}{n} \# \left\{ \bigcup_{j > k} \Pi_p(j) \cap [n] \right\} < \epsilon$$

From this identity, we deduce

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{n} \# \{ \Pi_t(i) \cap [n] \} &\leq \sum_{\substack{j \in \Pi_{p,t}(i) \\ j \leq k}} \limsup_{n \rightarrow \infty} \frac{1}{n} \# \{ \Pi_p(j) \cap [n] \} + \limsup_{n \rightarrow \infty} \frac{1}{n} \# \left\{ \bigcup_{\substack{j \in \Pi_{p,t}(i) \\ j > k}} \Pi_p(j) \cap [n] \right\} \\ &\leq \sum_{\substack{j \in \Pi_{p,t}(i) \\ j \leq k}} |\Pi_p(j)| + \epsilon \end{aligned}$$

and similarly

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{1}{n} \# \{ \Pi_t(i) \cap [n] \} &\geq \sum_{j \in \Pi_{p,t}(i), j \leq k} \liminf_{n \rightarrow \infty} \frac{1}{n} \# \{ \Pi_p(j) \cap [n] \} \\ &\geq \sum_{j \in \Pi_{p,t}(i), j \leq k} |\Pi_p(j)| \end{aligned}$$

This implies that $\Pi_t(i)$ admits an asymptotic frequency. The same kind of argument entails the existence of the asymptotic frequency of $\Pi_{t-}(i)$. This holds for all $i \geq 1$ and $t \geq 0$ simultaneously, almost surely. The càdlàg property of the frequencies derives from the preceding arguments. $\square$

8.2 Proof of Lemma 3.11

First, one has

$$\mathbb{P}[\hat{\Pi}_{r,t} = \text{Coag}(\hat{\Pi}_{s,t}, \hat{\Pi}_{r,s}), \forall r < s < t \in \mathbb{Q}^3] = 1 \quad (45)$$

To end the proof, it suffices to show that almost surely for every integer $n \geq 1$, the sequence

$$(\# \{ i \in [m] : \hat{\Pi}_{\frac{i-1}{m}, \frac{i}{m}}^{[n]} \neq 0_{[n]} \})_{m \geq 1}$$

is bounded. This is trivially the case for a stochastic flow of partitions $\hat{\Pi}^{\mathcal{P}}$ defined from a Λ lookdown graph $\mathcal{P}$. Using Equation (45) and the fact that the finite-dimensional marginals of $\hat{\Pi}$ have the same distribution as those of $\hat{\Pi}^{\mathcal{P}}$, we deduce that almost surely the sequence is bounded. $\square$

8.3 Proof of Lemma 3.12

We work on the event $\Omega_{\hat{\Pi}}$ throughout this proof, and recall the cocycle property, the left and right regularity properties verified by the flow restricted to its rational marginals. Fix $s \in \mathbb{Q}$ and let us prove the existence of the two limits asserted in the lemma. Since $(\hat{\Pi}_{s,v}, v \in [s, \infty) \cap \mathbb{Q})$ is the restriction of a càdlàg modification of $(\hat{\Pi}_{s,v}, v \in [s, \infty))$ to its rational marginals, the first limit is well-defined. Fix $t \in \mathbb{Q}$ and $n \in \mathbb{N}$, and let us prove the existence of the second limit. From the right regularity, we deduce that there exists $\epsilon > 0$ (that depends on $\omega \in \Omega_{\hat{\Pi}}$) such that $s + \epsilon < t$ and for all rationals $p < q \in (s, s + \epsilon)^2$, $\hat{\Pi}_{p,q}^{[n]} = 0_{[n]}$. Using the cocycle property, we easily deduce that $p \mapsto \hat{\Pi}_{p,t}^{[n]}$ is constant on $(s, s + \epsilon) \cap \mathbb{Q}$ and the limit exists.

For any $r < s < t$ such that at least s is rational or r and t are rational, we have

$$\tilde{\Pi}_{r,t} = \text{Coag}(\tilde{\Pi}_{s,t}, \tilde{\Pi}_{r,s})$$

This is due to the continuity of the coagulation operator (see Subsection 2.1) by passing to the limit on the cocycle property applied to the rational marginals.

Finally, suppose that $s, t \notin \mathbb{Q}$. To verify that our definition of $\tilde{\Pi}_{s,t}$ makes sense, we need to show that $\text{Coag}(\tilde{\Pi}_{q,t}, \tilde{\Pi}_{s,q})$ does not depend on the value $q \in (s, t) \cap \mathbb{Q}$. Consider two such values $q, q' \in (s, t) \cap \mathbb{Q}$, suppose that $q < q'$ and use Equation (5) to obtain

$$\begin{aligned} \text{Coag}(\tilde{\Pi}_{q',t}, \tilde{\Pi}_{s,q'}) &= \text{Coag}(\tilde{\Pi}_{q',t}, \text{Coag}(\tilde{\Pi}_{q,q'}, \tilde{\Pi}_{s,q})) = \text{Coag}(\text{Coag}(\tilde{\Pi}_{q',t}, \tilde{\Pi}_{q,q'}), \tilde{\Pi}_{s,q}) \\ &= \text{Coag}(\tilde{\Pi}_{q,t}, \tilde{\Pi}_{s,q}) \end{aligned}$$

Thus, the definition of $\tilde{\Pi}_{s,t}$ does not depend on $q \in (s, t)$.

Finally, consider three irrational $r < s < t$, and two rational values q, q' such that $q \in (r, s)$ and $q' \in (s, t)$.

$$\begin{aligned} \text{Coag}(\tilde{\Pi}_{s,t}, \tilde{\Pi}_{r,s}) &= \text{Coag}(\text{Coag}(\tilde{\Pi}_{q',t}, \tilde{\Pi}_{s,q'}), \text{Coag}(\tilde{\Pi}_{q,s}, \tilde{\Pi}_{r,q})) \\ &= \text{Coag}(\tilde{\Pi}_{q',t}, \text{Coag}(\tilde{\Pi}_{s,q'}, \text{Coag}(\tilde{\Pi}_{q,s}, \tilde{\Pi}_{r,q}))) = \text{Coag}(\tilde{\Pi}_{q',t}, \tilde{\Pi}_{r,q'}) = \tilde{\Pi}_{r,t} \end{aligned}$$

This concludes the proof. $\square$

8.4 Proof of Proposition 3.13

By definition, for every rational numbers $s \leq t$, $\tilde{\Pi}_{s,t} = \hat{\Pi}_{s,t}$ on the event $\Omega_{\hat{\Pi}}$, so it holds a.s. Suppose that $s \in \mathbb{Q}$. On the event $\Omega_{\hat{\Pi}}$, $(\tilde{\Pi}_{s,t}, t \in [s, \infty))$ coincides with a càdlàg modification of $(\hat{\Pi}_{s,t}, t \in [s, \infty))$. Therefore, for every $t > s$, a.s. $\tilde{\Pi}_{s,t} = \hat{\Pi}_{s,t}$.

Now suppose that $s \in \mathbb{R} \setminus \mathbb{Q}$, $t > s$ and fix $n \in \mathbb{N}$. We have for all $q \in (s, t) \cap \mathbb{Q}$

$$\mathbb{P}(\hat{\Pi}_{s,t}^{[n]} = \hat{\Pi}_{s,t}^{[n]}) \geq \mathbb{P}(\tilde{\Pi}_{q,t}^{[n]} = \hat{\Pi}_{q,t}^{[n]}; \hat{\Pi}_{s,t}^{[n]} = \text{Coag}(\hat{\Pi}_{q,t}^{[n]}, \hat{\Pi}_{s,q}^{[n]}; \hat{\Pi}_{s,q}^{[n]} = 0_{[n]})$$

As $q \downarrow s$, $\mathbb{P}(\hat{\Pi}_{s,q}^{[n]} = 0_{[n]}) \rightarrow 1$ by definition of a stochastic flow of partitions. The cocycle property of a stochastic flow of partitions together with the almost sure identity $\tilde{\Pi}_{q,t} = \hat{\Pi}_{q,t}$ we already proved, ensures that the probability of the event on the r.h.s. tends to 1 as $q \downarrow s$. Thus $\tilde{\Pi}_{s,t} = \hat{\Pi}_{s,t}$ almost surely. Finally, when $t = s$ we know that $\hat{\Pi}_{s,s} = 0_{[\infty]}$ almost surely by definition. Therefore, $\tilde{\Pi}$ is a modification of $\hat{\Pi}$.

We need to verify that for all $\omega \in \Omega_{\hat{\Pi}}$, $\tilde{\Pi}(\omega)$ is a deterministic flow of partitions. The cocycle property was proved in the preceding lemma. Let us show the left and right regularities. Fix $s \in \mathbb{R}$ and $n \in \mathbb{N}$. Recall that there exists $\epsilon > 0$ such that for all rational $p < q \in [(s, s + \epsilon) \cap \mathbb{Q}]^2$, $\tilde{\Pi}_{p,q}^{[n]} = 0_{[n]}$. Letting $p \downarrow s$, we get $\tilde{\Pi}_{s,q}^{[n]} = 0_{[n]}$ for all $q \in (s, s + \epsilon) \cap \mathbb{Q}$. Similarly for all $r \in (s, s + \epsilon)$, we have $\tilde{\Pi}_{q,r}^{[n]} = 0_{[n]}$ as soon as $q \in (s, r)$. Using the fact that $\tilde{\Pi}_{s,r}^{[n]} = \text{Coag}(\tilde{\Pi}_{q,r}^{[n]}, \tilde{\Pi}_{s,q}^{[n]})$ we get that $\tilde{\Pi}_{s,r}^{[n]} = 0_{[n]}$ for all $r \in (s, s + \epsilon)$. This in turn implies that $\tilde{\Pi}_{s,r}^{[n]} \rightarrow \tilde{\Pi}_{s,s}^{[n]}$ as $r \downarrow s$ and the right regularity is proved. The left regularity is obtained similarly. $\square$

8.5 Proof of Proposition 4.7

To alleviate notation, we prove the proposition for $s = 0$. Let $\mathcal{M}_f(a)$ be the set of finite measures on $[0, a]$, equipped with its weak topology. Consider a $\mathcal{M}_f(1)$ -valued process $(m_t, t \geq 0)$ associated with

Neveu continuous state branching process (CSBP in short) as defined in [9]. Therefore $(Z_t(x), t \geq 0) := (m_t([0, x]), t \geq 0)$ is a Neveu CSBP started from x , for each $x \in [0, 1]$. We define for all $t \geq 0$

$$\rho_t(dx) := \frac{m_t(dx)}{m_t([0, 1])}$$

$(\rho_t, t \geq 0)$ is a Λ Fleming-Viot process where $\Lambda(dx) = dx$ (see [9] for a proof of this result). Hence, we will prove the proposition by considering the process $(m_t, t \geq 0)$.

For all $t > 0$, the distribution function $(S_t(x), x \in [0, 1])$ of m_t is a subordinator without drift whose Lévy measure has an infinite mass (see Section 3 in [5]). Denote by $\mathcal{U}$ the set of ancestral types of this measure-valued process, that is

$$\mathcal{U} := \{x \in [0, 1] : \exists t > 0, m_t(x) > 0\} \quad (46)$$

Lemma 8.1 *The set $\mathcal{U}$ is a countable subset of $[0, 1]$. For each $u \in \mathcal{U}$, for all $t > 0$, $m_t(u) > 0$.*

Proof Since the distribution function of m_t has no drift part, the set of atoms of m_{t+s} is included in the set of atoms of m_t for all $t, s > 0$. Moreover, if $m_t(u) > 0$ for a given time $t > 0$ and a given point $u \in [0, 1]$, then the process $(m_{t+s}(u), s \geq 0)$ is a Neveu CSBP started from $m_t(u)$, which is independent of $(m_{t+s}([0, 1] \setminus \{u\}), s \geq 0)$. Since a Neveu CSBP does not get extinct in finite time almost surely, we deduce that an ancestral type has a positive mass at any time. $\square$

We now prove that we can order the set $\mathcal{U}$ according to the formula given in the statement of the proposition. We restrict our proof to the first two Eves e^1 and e^2 , since the general proof will then be a simple consequence. For each $n \geq 1$, consider the intervals

$$[0, \frac{1}{n}), [\frac{1}{n}, \frac{2}{n}), \dots, [\frac{n-1}{n}, 1]$$

The processes obtained by restricting $(m_t, t \geq 0)$ to each of these intervals are themselves measure-valued processes associated with Neveu branching mechanism, and are independent thanks to the branching property. Additionally, each of them fixes on a primitive Eve, say $e(i, n)$, for the primitive Eve corresponding to the process

$$(m_t(\cdot \cap [\frac{i-1}{n}, \frac{i}{n})), t \geq 0)$$

for every $i \in [n]$. Obviously $\{e(i, n), i \in [n]\} \subset \mathcal{U}$ for every $n \geq 1$. Moreover, one can prove that for two distinct integers $i \neq j \in [n]$, we have

$$\frac{m_t(\{e(i, n)\})}{m_t(\{e(j, n)\})} \xrightarrow{t \uparrow \infty} \{0, \infty\}$$

This ensures the existence of two integers $i_1^n, i_2^n \in [n]$ such that

$$\begin{aligned} \frac{m_t(\{e(i_1^n, n)\})}{m_t(\{e(j, n)\})} &\xrightarrow{t \uparrow \infty} \infty \text{ for all } j \neq i_1^n \\ \frac{m_t(\{e(i_2^n, n)\})}{m_t(\{e(j, n)\})} &\xrightarrow{t \uparrow \infty} \infty \text{ for all } j \neq i_1^n, i_2^n \end{aligned}$$

We claim that both sequences $(e(i_1^n, n))_{n \geq 1}$ and $(e(i_2^n, n))_{n \geq 1}$ are eventually constant almost surely. This is clear for the first one, since for all $n \geq 1$, $e(i_1^n, n) = \mathbf{e}^1$ that is the primitive Eve of the entire population. Concerning the second, suppose that there exists an event C of probability greater than 0 and a random increasing sequence $(n_k)_{k \geq 1}$ of integers such that on C , $e(i_2^{n_k}, n_k) \neq e(i_2^{n_k-1}, n_k - 1)$ for every $k \geq 1$. A compatibility argument on the restrictions of the process m ensures that for every $k \geq 1$

$$e(i_2^{n_k}, n_k) \in \left[\frac{i_1^{n_k-1} - 1}{n_k - 1}, \frac{i_1^{n_k-1}}{n_k - 1} \right)$$

and therefore $|\mathbf{e}^1 - e(i_2^{n_k}, n_k)| \rightarrow 0$ as $k \uparrow \infty$ on event C . But this is not possible since for every $n \geq 1$, the pair (i_1^n, i_2^n) is uniformly distributed among the pair of distinct integers in $[n]$. Hence $|\mathbf{e}^1 - e(i_2^{n_k}, n_k)| \rightarrow 0$ has probability 0. Finally the claimed property is verified.

We then denote by $\mathbf{e}^2$ the last value taken by the sequence $(e(i_2^n, n))_{n \geq 1}$. From its definition, we obtain the convergence

$$\frac{m_t(\{\mathbf{e}^2\})}{m_t([0, 1] \setminus \{\mathbf{e}^1\})} \xrightarrow[t \uparrow \infty]{} 1$$

The assertion of the proposition follows. $\square$

8.6 Proof of Lemma 5.4

Suppose *a*). Then we know that $B_t \xrightarrow{(d)} Id$ as $t \downarrow 0$ from the Continuity Lemma 1, in [6]. Since the limit is a continuous function, we have for every $0 \leq x \leq 1$, $B_t(x) \xrightarrow{(d)} x$. The limit being deterministic, the convergence also holds in probability $B_t(x) \xrightarrow{(\mathbb{P})} x$.

Fix $\epsilon > 0$. Denote by $[x]$ the integer part of any real x . There exists $t_0 > 0$ such that for every $t \in (0, t_0)$ and $k \in [\lfloor 1/\epsilon \rfloor]$

$$\mathbb{P}(|B_t(k\epsilon) - k\epsilon| > \epsilon) < \frac{\epsilon}{2^k}$$

From the monotonicity of B_t , we get $\mathbb{P}(\|B_t - Id\|_\infty > 2\epsilon) < 2\epsilon$. Hence, $B_t \xrightarrow{(\mathbb{P})} Id$.

Suppose *b*). Fix $n \geq 1$ and $\epsilon > 0$, we will prove there exists $t_0 > 0$ such that for all $t \in (0, t_0)$

$$\mathbb{P}(d_{\mathcal{D}}(\pi(B_t), \pi(Id)) < 2^{-n}) > 1 - 2\epsilon$$

There exists $p \in \mathbb{N}$ such that

$$\mathbb{P}(\{\exists i, j \text{ s.t. } 1 \leq i < j \leq n \text{ and } |V_i - V_j| < \frac{2}{p}\}) < \epsilon \quad (47)$$

Moreover, there exists $t_0 > 0$ such that for all $t \in (0, t_0)$

$$\mathbb{P}(\bigcap_{0 \leq k \leq p} \{B_t(\frac{k}{p}) \in [\frac{k}{p} - \frac{1}{3p}, \frac{k}{p} + \frac{1}{3p}]\}) > 1 - \epsilon \quad (48)$$

The monotonicity of B_t^{-1} , and the two previous equations ensure that

$$\mathbb{P}(\{(B_t^{-1}(V_1), \dots, B_t^{-1}(V_n)) \text{ are all distinct}\}) > 1 - 2\epsilon$$

Thus, we obtain a)

□

Acknowledgements. This is part of my PhD thesis. I would like to thank my supervisors: Julien Berestycki for his very useful advice throughout this work and Amaury Lambert for his careful reading of a first version of this paper.

References

- [1] J. BERESTYCKI AND N. BERESTYCKI, *Kingman's coalescent and Brownian motion*, Alea, 6 (2009), pp. 239–259.
- [2] J. BERESTYCKI, N. BERESTYCKI, AND J. SCHWEINSBERG, *Beta-coalescents and continuous stable random trees.*, Ann. Probab., 35 (2007), pp. 1835–1887.
- [3] J. BERESTYCKI, N. BERESTYCKI, AND J. SCHWEINSBERG, *Small-time behavior of beta coalescents.*, Annales de l'Institut Henri Poincaré (B) Probability and Statistics, 44 (2008), pp. 214 – 238.
- [4] J. BERTOIN, *Random fragmentation and coagulation processes*, Cambridge University Press, Cambridge, 2006.
- [5] J. BERTOIN AND J.-F. LE GALL, *The Bolthausen-Sznitman coalescent and the genealogy of continuous-state branching processes*, Probability Theory and Related Fields, 117 (2000), pp. 249–266.
- [6] ———, *Stochastic flows associated to coalescent processes*, Probability Theory and Related Fields, 126 (2003), pp. 261–288.
- [7] ———, *Stochastic flows associated to coalescent processes II: Stochastic differential equations*, Annales de l'Institut Henri Poincaré (B) Probability and Statistics, 41 (2005), pp. 307 – 333.
- [8] ———, *Stochastic flows associated to coalescent processes. III. Limit theorems*, Illinois J. Math., 50 (2006), pp. 147–181 (electronic).
- [9] M. BIRKNER, J. BLATH, M. CAPALDO, A. M. ETHERIDGE, M. MÖHLE, J. SCHWEINSBERG, AND A. WAKOLBINGER, *Alpha-stable branching and beta-coalescents*, Electronic Journal of Probability, 10 (2005), pp. 303–325.
- [10] M. BIRKNER, J. BLATH, M. MOEHLE, M. STEINRUECKEN, AND J. TAMS, *A modified look-down construction for the Xi-Fleming-Viot process with mutation and populations with recurrent bottlenecks*, Alea, 6 (2009), pp. 25–61.
- [11] P. DONNELLY AND T. G. KURTZ, *A countable representation of the Fleming-Viot measure-valued diffusion.*, Ann. Probab., 24 (1996), pp. 698–742.

- [12] ———, *Particle representations for measure-valued population models.*, Ann. Probab., 27 (1999), pp. 166–205.
- [13] T. DUQUESNE AND J.-F. LE GALL, *Random Trees, Levy Processes and Spatial Branching Processes*, Astérisque, (2005).
- [14] C. FOUCART, *Generalized Fleming-Viot processes with immigration via stochastic flows of partitions*, ArXiv e-prints, (2011).
- [15] O. KALLENBERG, *Canonical representations and convergence criteria for processes with interchangeable increments*, Probability Theory and Related Fields, 27 (1973), pp. 23–36. 10.1007/BF00736005.
- [16] P. PFAFFELHUBER AND A. WAKOLBINGER, *The process of most recent common ancestors in an evolving coalescent*, Stochastic Processes and their Applications, 116 (2006), pp. 1836 – 1859.
- [17] J. PITMAN, *Coalescents with multiple collisions*, Ann. Probab, 27 (1999), pp. 1870–1902.
- [18] S. SAGITOV, *The general coalescent with asynchronous mergers of ancestor lines.*, J.Appl.Prob., 36 (1999), pp. 1116–1125.
- [19] J. SCHWEINSBERG, *A necessary and sufficient condition for the Lambda-coalescent to come down from infinity.*, Electronic Communications in Probability, 5 (2000), pp. 1–11.